

# Responses of Microbial Communities to Single-Walled Carbon Nanotubes in Phenol Wastewater Treatment Systems

Yuanyuan Qu,<sup>\*,†,‡,||</sup> Qiao Ma,<sup>†,‡,||</sup> Jie Deng,<sup>‡,||</sup> Wenli Shen,<sup>†</sup> Xuwang Zhang,<sup>†</sup> Zhili He,<sup>‡</sup> Joy D. Van Nostrand,<sup>‡</sup> Jiti Zhou,<sup>†</sup> and Jizhong Zhou<sup>‡,§,⊥</sup>

<sup>†</sup>Key Laboratory of Industrial Ecology and Environmental Engineering (Ministry of Education), School of Environmental Science and Technology, Dalian University of Technology, Dalian 116024, China


<sup>‡</sup>Institute for Environmental Genomics (IEG), Department of Microbiology and Plant Biology, University of Oklahoma, Norman, Oklahoma 73019, United States

<sup>§</sup>Earth Sciences Division, Lawrence Berkeley National Laboratory, Berkeley, California 94720, United States

<sup>⊥</sup>State Key Joint Laboratory of Environment Simulation and Pollution Control, School of Environment, Tsinghua University, Beijing 100084, China

## Supporting Information

**ABSTRACT:** The expanding use of single-walled carbon nanotubes (SWCNTs) raises environmental concerns. Wastewater treatment systems are potential recipients of SWCNTs containing influent, yet the impacts of SWCNTs on these systems are poorly documented. In this study, the microbial responses to SWCNTs in simulated phenol wastewater treatment systems were investigated. The phenol removal rates were improved in all SWCNTs-treated sequencing batch reactors during the first 20 days, but when facing higher phenol concentration (1000 mg/L) after 60 days, reactors with the highest concentration (3.5 g/L) of SWCNTs exhibited a notably decreased phenol removal capacity. Cell viability tests, scanning electron microscopy analysis and DNA leakage data suggested that SWCNTs protected microbes from inactivation, possibly by producing more bound extracellular polymeric substances (EPS), which could create a protective barrier for the microbes. Illumina sequencing of 16S rRNA gene amplicons revealed that the bacterial diversity did not change significantly except for a minor reduction after the immediate addition of SWCNTs. Bacterial community structure significantly shifted after SWCNTs addition and did not recover afterward. *Zoogloea* increased significantly upon SWCNTs shocking. At the final stage, *Rudaea* and *Mobilicoccus* increased, while *Burkholderia*, *Singulisphaera*, *Labrys* and *Mucilaginibacter* decreased notably. The shifts of these dominant genera may be associated with altered sludge settling, aromatic degradation and EPS production. This study suggested that SWCNTs exerted protective rather than cytotoxic effects on sludge microbes of phenol wastewater treatment systems and they affected the bacterial community structure and diversity at test concentrations. These findings provide new insights into our understanding of the potential effects of SWCNTs on wastewater treatment processes.



## INTRODUCTION

Carbon nanotubes (CNTs) are among the most promising engineering nanomaterials due to their unique physicochemical properties compared to bulk materials. They have been incorporated into a diverse array of commercial products such as pharmaceuticals, optical devices, cosmetics, electronics and antimicrobial coatings.<sup>1,2</sup> With the exponential increase in manufacturing and application of CNTs in nanotechnology, they will inevitably enter various environmental matrices.<sup>3-6</sup> However, the potential environmental risks are poorly understood and a comprehensive investigation is needed.

Cytotoxicity of CNTs to microbes has been demonstrated using pure microbial strains.<sup>7-13</sup> The underlying molecular mechanisms were proposed to be the synergistic impacts of cell

membrane perturbation and oxidative stress. However, evidence has shown that CNT toxicity to pure microbial strains is a poor predictor of toxicity to microbial communities.<sup>14,15</sup> A number of studies have therefore investigated the effects of CNTs on microbial communities in aquatic and soil environments.<sup>16-18</sup> For instance, Chung et al. reported that high concentrations of CNTs significantly lowered biomass and some enzyme activities of microbial communities from an urban soil,<sup>19,20</sup> whereas Shrestha et al.

Received: October 31, 2014

Revised: March 2, 2015

Accepted: March 9, 2015

Published: March 9, 2015

suggested that soil respiration and extracellular enzyme activities were not significantly affected in a sandy loam soil even in the presence of extremely high concentrations of multiwalled carbon nanotubes (MWCNTs).<sup>21</sup> CNTs were also reported to affect microbial degradation of aromatic pollutants including 2,4-dichlorophenol and phenanthrene.<sup>22,23</sup> Therefore, it is necessary to investigate the responses of microbial communities to CNTs from various environments.

Wastewater treatment plants (WWTPs), as important receptors and the sink of waste streams, are among the most probable CNT recipients of industrial and domestic effluents.<sup>24</sup> Moreover, CNTs are being tailored to work as catalyst supports, composite filters, adsorbents and antimicrobial agents in wastewater treatment processes, increasing the likelihood of contact with microbial communities in activated sludges.<sup>2,25</sup> Consequently, they may pose a risk to the microbial populations and their associated functions. In this respect, the effects of other nanoparticles (Cu-, Zn-, Ag-, Al<sub>2</sub>O<sub>3</sub>-, ZnO-, TiO<sub>2</sub>-, SiO<sub>2</sub>- nanoparticles), especially Ag-nanoparticles, have been extensively studied and most negatively affected waste removal efficacy.<sup>26–28</sup> Only a few studies have examined the effects of CNTs on wastewater microbial communities.<sup>29–32</sup> Luongo and Zhang found that MWCNTs inhibited the activated sludge respiration in a dose-dependent manner within 3 h,<sup>30</sup> and that single-walled carbon nanotubes (SWCNTs) altered the structure of the bacterial communities in the sludge systems within 5 h based on automated ribosomal intergenic spacer analysis (ARISA).<sup>29</sup> Very recently, two studies also revealed that CNTs affected bacterial community structure of activated sludge and its corresponding functions including methane production, nitrogen and phosphorus removal.<sup>33,34</sup> Notwithstanding, the ecological effects of CNTs on activated sludge system and microbial survival are far from clear.

Recently, reports have demonstrated that CNTs altered the composition and structure of microbial communities based on results from several culture-independent technologies including ARISA, PCR-denaturing gradient gel electrophoresis (DGGE), multiplex-terminal restriction fragment length polymorphism (M-TRFLP) and clone library analyses.<sup>18,29,35</sup> However, these methods only allowed identification of microbial populations at rather coarse taxonomic levels. Current advances in high-throughput sequencing techniques have not only increased sequencing depth at a lower cost but also provided higher taxonomic resolution.<sup>36–38</sup> Especially, the Illumina MiSeq platform, which is able to achieve comparable or greater sequencing depth than related pyrosequencing, has been widely used to examine the phylogenetic/taxonomic diversity, composition and structure of microbial communities from a variety of environments.<sup>21,39</sup>

In this study, we aimed to (1) explore the impacts of SWCNTs on pollutant removal efficiency in activated sludge systems, (2) investigate the microbial survival and cytotoxic mechanism, (3) monitor diversity and structure shifts of microbial communities and (4) identify the dominant microorganisms susceptible to SWCNTs. To achieve these goals, sequencing batch reactors (SBRs) were constructed for treating phenol containing wastewaters that were dosed with 0.5, 1.5, 2.5 and 3.5 g/L SWCNTs. Results demonstrated that SWCNTs played protective roles for sludge microbes, and in the meantime, changed the structure of sludge bacterial communities.

## ■ MATERIALS AND METHODS

**SWCNTs and Activated Sludge.** Commercially available SWCNTs (>95%) were purchased from Shenzhen Nanotech Port Co., Ltd. (Shenzhen, China) and were suspended in distilled water using ultrasonic treatment for 30 min to obtain a better dispersion. More details regarding the SWCNTs have been described previously.<sup>40</sup> Activated sludge was gathered from the secondary sedimentation tank of Chunliu River WWTP (Dalian, China).

**Experimental Design.** The reactors were 65 cm tall with an internal diameter of 8 cm and a working volume of 2.5 L. Fine air bubbles for aeration were supplied through an air pump at the reactor bottom with an airflow rate of 0.4 L/min. The synthetic wastewater consisted of 20 mg of KH<sub>2</sub>PO<sub>4</sub>/L, 90 mg of NH<sub>4</sub>Cl/L, 10 mg of NaCl/L, 12.5 mg of MgSO<sub>4</sub>·7H<sub>2</sub>O/L, 12 mg of CaCl<sub>2</sub>/L, 10 mg of FeSO<sub>4</sub>·7H<sub>2</sub>O/L, 785 mg of glucose/L and 200 mg phenol/L. The SBRs (*n* = 15) were seeded with activated sludge (2.57 g/L, dry weight at 105 °C), and domesticated with the synthetic wastewater. The SBRs were operated under identical conditions. Each cycle of SBR was operated for 24 h, including 2 h of fill, 18 h of aeration, 2 h of settling, and 2 h of decant. Effluent was discharged with a volumetric exchange ratio of 50%. After 15 days of domestication, the SBRs were divided into 5 groups, each receiving different concentrations of SWCNT (g/L): 0 (G1), 0.5 (G2), 1.5 (G3), 2.5 (G4) and 3.5 (G5). Each group contained three replicates. During the 2 month operation period, phenol concentrations of the influent and effluent were monitored daily, and the concentrations of mixed liquor suspended solid (MLSS), the sludge volume after 0.5 h of settling (SV<sub>30</sub>), pH and dissolved oxygen (DO) were measured every other day. Because the system was operated in good conditions, and there was barely any sludge discharge during the whole process, the MLSS was relatively stable during the process (Figure S1 of the Supporting Information), thus the sludge retention time was not considered herein. The influent phenol concentration was increased to 1000 mg/L on Day 61 to compare the robustness of the control reactors with the SWCNTs-treated ones. Phenol removal was monitored on Day 1, 20, 40, 60, 61 and 67 (the last day of operation).

**Analytical Methods.** Because activated sludge used in this study was gathered from a municipal WWTP and aerated immediately, which was transparent and colorless after centrifugation, the concentration of phenol was measured directly using a UV-vis spectrophotometer (V-560, JASCO, Japan). MLSS and SV<sub>30</sub> were determined according to standard methods. The pH and DO were measured using a pH meter (S20, Mettler-Toledo, Switzerland) and a DO meter (FLX310, Flow Electronic, China), respectively. Scanning electron microscopy (SEM) images of the SWCNTs and activated sludges were recorded using field emission scanning electron microscopy (FE-SEM, KYKY-1000B, KYKY Technology, China). For the detection of live/dead cells, the 2-(4-amidinophenyl)-6-indolecarbamide dihydrochloride (DAPI, Beyotime, China) and propidium iodide (PI, Beyotime, China) staining assays were performed according to the manufacturer's instructions. The efflux DNA was determined by fluorescence spectroscopy (Hitachi, Japan) using DAPI as the fluorescent dye (excitation 364, emission 454 nm) after filtration through a 0.22 μm membrane. The bound extracellular polymeric substance (EPS) of activated sludge was extracted using ethylenediaminetetraacetic acid. Concentrations of protein

and carbohydrate were measured using the Lowry and anthrone methods, respectively.<sup>41</sup>

**DNA Extraction, PCR Amplification and Sequencing.** Activated sludge samples were collected at Day 1, 20, 40, 60 and 67 before sludge settling, and the genomic DNA was extracted using a protocol based on Purkhold et al.<sup>42,43</sup> DNA concentration was measured with Pico Green assays using a FLUOstar OPTIMA fluorescence plate reader (BMG Labtech, Germany). For high-throughput sequencing, the primers 515F (5'-GTG CCAG CMGC CGCG GTAA-3') and 806R (5'-GGAC TACH VGGG TWTC TAAT-3') were used to amplify the V4 region of the 16S rRNA gene.<sup>44</sup> PCR was conducted in a 25  $\mu$ L mixture containing 0.1  $\mu$ L of AccuPrime High Fidelity Taq Polymerase (Invitrogen, USA), 2.5  $\mu$ L of 10 $\times$  AccuPrime PCR buffer II, 1  $\mu$ L of each primer (10  $\mu$ M), 1  $\mu$ L of template DNA and 19.4  $\mu$ L of nuclease-free water under the following conditions: 94  $^{\circ}$ C for 1 min; 35 cycles of 94  $^{\circ}$ C for 20 s, 53  $^{\circ}$ C for 25 s, and 68  $^{\circ}$ C for 45 s; final extension at 68  $^{\circ}$ C for 10 min. Each sample was amplified in triplicate. PCR products were pooled, purified through QIAquick Gel Extraction Kit (Qiagen), and quantified by Pico Green analysis. The 16S rRNA high-throughput sequencing was conducted on Illumina MiSeq platform at the Institute for Environmental Genomics, University of Oklahoma.


**Sequencing Data Analysis of 16S rRNA Gene Amplicons.** After sequencing, PhiX sequences were removed and primers were trimmed (mismatch 1.5), and the paired-end reads were joined using the Flash program (phredOffset 33, standard deviation of fragment lengths 20).<sup>45</sup> Sequences containing ambiguous reads (N) and reads shorter than 240 bp were removed.<sup>46</sup> The resulting sequences were screened for Chimeras using UCHIME.<sup>47</sup> Operational taxonomic units (OTUs) were categorized using CD-HIT at a 97% sequence similarity threshold,<sup>48</sup> and the taxonomic assignment of OTUs was performed by RDP classifier with 50% confidence.<sup>49</sup> The above processes were performed through a pipeline (<http://zhoulab5.rccc.ou.edu/>) (not published). Detrended correspondence analysis (DCA) and correlation tests were calculated using R v2.15.1 (<http://www.r-project.org/>). Three non-parametric tests, including multiple response permutation procedure (MRPP), permutational multivariate analysis of variance (Adonis) and analysis of similarity (ANOSIM) were performed to test dissimilarity among treatment groups based on bray-cutis distance index (<http://ieg.ou.edu/>).

## RESULTS AND DISCUSSION

**Characterization of SWCNTs.** Cytotoxicity of CNTs was closely related to the physicochemical properties. The pristine SWCNTs used here were 5–16  $\mu$ m in length, and less than 2 nm in diameter, with a surface area of 500–700 m<sup>2</sup>/g. SEM and transmission electron microscopy (TEM) micrographs were obtained to confirm the manufacturer's description.<sup>40</sup> The impact of low metal impurities (carbon content > 95%) was not considered since it showed insignificant antibacterial activity on soil and activated sludge communities.<sup>15,29,35</sup> With the increasing manufacturing and application of CNTs, more and more CNTs will inevitably enter WWTPs via CNT production facility's release, manufacturing and disposal of CNT containing products.<sup>4,5,50,51</sup> CNTs released from composites such as sports equipment, tires, electronics, etc. have been reported elsewhere.<sup>51</sup> CNTs may agglomerate and accumulate in activated sludge due to their high hydrophobic and biodegradation-resistant characteristics.<sup>5,34</sup> In previous reports, low concen-


trations of CNTs have been proven to exert no or little impact on microbial and enzyme activities of soils and activated sludge.<sup>19,33,35</sup> Our preliminary experiments also showed that the microbial communities were not changed with 0.1 g/L SWCNT, whereas a noticeable shift was observed in the 2.0 g/L SWCNT group using PCR-DGGE analysis (data not shown). Therefore, the concentrations of SWCNTs used in the present study were set at a relative high level from 0.5 to 3.5 g/L.

**Effects of SWCNTs on Reactor Performances.** Acute and chronic influences of SWCNTs on the phenol wastewater treatments were mimicked over a course of 2 months of operation. Phenol was almost completely removed after one cycle of operation in all reactors (Figure 1). In the first 20 days,



**Figure 1.** Phenol removal performances of each group during the 67 day operation process. Each group was performed in triplicate with different concentrations of SWCNTs (g/L): 0 (G1), 0.5 (G2), 1.5 (G3), 2.5 (G4) and 3.5 (G5).

phenol removal rates were SWCNT-dose-dependent with the order of G5 > G4 > G3 > G2 > G1 (Figure S2 of the Supporting Information). On Day 1, the removal rate reached  $96.1 \pm 2.4\%$  within 2 h in G5, whereas it took 8 h for G1 to reach a similar removal rate, indicating a positive influence of SWCNTs on phenol removal in this system. After 40 days of operation, all groups reached similar phenol removal rates (Figure S2 of the Supporting Information). Therefore, despite the potential toxicity, SWCNTs appeared to exhibit positive effects on sludge microbial communities, especially during the early stages of operation. SWCNTs have been widely used as adsorbents for heavy metals and aromatics removal by virtue of their unique properties.<sup>52–54</sup> Results from this study also indicated a dose-dependent relationship between phenol removal and SWCNTs, and 3.5 g/L SWCNTs could adsorb 58.9% phenol (initial concentration 180 mg/L) within 6 h (Table S1 of the Supporting Information). Thus, we initially speculated the elevated phenol removal rates on Day 1 and 20 were attributed to phenol adsorption by SWCNTs. However, when autoclaved sludge systems were used, phenol adsorption did not improve in SWCNTs-treated groups (Table S1 of the Supporting Information). It was previously proven that carbon-based nanomaterials could adsorb free EPS and improve EPS production, which could form a protective barrier for the microbes.<sup>12,31,34</sup> Therefore, we determined the bound protein and polysaccharide concentrations of the five groups (Figure S3 of the Supporting Information), which revealed higher bound EPS concentrations in SWCNTs-treated SBRs compared to the control group. Altogether, our results suggested that the potential protective mechanism of SWCNTs possibly resulted from the higher bound EPS production.



**Figure 2.** DAPI/PI staining results of each group on Day 1. Blue parts represent the active cells stained with DAPI and red parts represent the inactivated cells stained with PI.


**Table 1.**  $\alpha$ -Diversity of All the Groups at Each Sampling Time

group	Day 1					Day 20				
	G1	G2	G3	G4	G5	G1	G2	G3	G4	G5
Shannon index	3.85	3.61	3.56	3.48	3.48	2.66	2.60	2.54	2.50	2.81
evenness	0.61	0.57	0.56	0.56	0.56	0.49	0.46	0.44	0.43	0.48
Chao1	861	868	926	857	824	401	484	526	595	630
OTU	533	543	550	533	522	237	300	325	334	364
group	Day 40					Day 60				
	G1	G2	G3	G4	G5	G1	G2	G3	G4	G5
Shannon index	3.37	3.69	4.00	3.87	3.98	3.79	3.46	3.41	3.49	3.92
evenness	0.57	0.61	0.65	0.63	0.66	0.65	0.65	0.59	0.57	0.59
Chao1	553	655	742	695	665	502	540	654	596	663
OTU	379	405	474	482	418	340	348	387	369	435
group	Day 67									
	G1	G2	G3	G4	G5					
Shannon index	3.10	3.00	2.84	2.87	3.16					
evenness	0.64	0.61	0.59	0.59	0.65					
Chao1	578	559	565	567	579					
OTU	340	345	324	349	368					

The phenol concentration in the influent was increased to 1000 mg/L on Day 61 to investigate the robustness of the constructed systems. Over 94% of the phenol was removed within 14 h in all groups, with a slightly lower removal rate in G5 (Figure S2 of the Supporting Information). On Day 67, all five groups showed different removal performances. G1 had a similar removal rate compared to Day 61; G2, G3 and G4 showed higher removal rates, in which 96% phenol was removed within 10 h; G5 had a considerably lower removal rate, requiring over 18 h to achieve 88% phenol removal (Figure S2 of the Supporting Information). Therefore, different concentrations of SWCNTs had different effects on wastewater

treatment systems. At low concentrations, they improved treatment performance. Once past a threshold, i.e., 3.5 g/L in the present study, they could result in a negative impact after a longtime interaction.

To determine cell survival rates upon addition of SWCNTs, cell viability tests (DAPI/PI staining) was performed at each sampling time (Figure 2, Figure S4 of the Supporting Information). The percentage of inactivated cells in the control group was similar to G2 and was much higher than those in G3, G4 and G5. This phenomenon was consistent throughout the whole operation. Therefore, the addition of SWCNTs at relatively high concentrations apparently lowered phenol



**Figure 3.** DCA plots of all samples showing the relationships of microbial community structures among different groups at each sampling time (Day 1, 20, 40, 60 and 67). Symbols represent the samples from different groups. A distinct cluster can be defined from the samples of G1 based on the DCA ordination, which suggests the differences in microbial community structures between G1 and other four groups.

toxicity to cells. DNA leakage data also revealed a higher cell-free DNA intensity in G1, confirming the lower cell membrane damage rates in the SWCNT groups (Figure S5 of the Supporting Information). The results further confirmed the positive influences of SWCNTs on the vitality of the microbial communities, which were also consistent with the EPS data. Three major cytotoxic mechanisms of CNTs have been recognized in previous studies, i.e., generation of oxidative stress, release of certain impurities and physical perturbation.<sup>33,34,55</sup> The reduced cell death and high purity of SWCNTs in the present study implied oxidative stress and metal purities should not be significant. SEM images showed a prevalence of

aggregates of SWCNTs, largely reducing their direct contact with microbial cells (Figure S6 of the Supporting Information). Only a few morphologically changed cells intertwined with SWCNTs, indicating limited physical toxicity of SWCNTs.<sup>15,29</sup> Meanwhile, the bound EPS concentrations in SWCNTs-treated groups were higher than those of the control group, further reducing the possibility of SWCNT physical toxicity. Therefore, the cytotoxicity of SWCNTs on sludge microbial communities was very limited and negligible. However, our results also suggested that G5 displayed a decreased phenol removal capability after phenol shock on Day 61, contrary to the results derived from microscopic observations. The distinct phenol

degrading capability could be primarily, if not completely, due to community shifts resulting from SWCNTs addition.

**Shifts of Bacterial Community Diversity and Structure.** The sludge samples from Day 1, 20, 40, 60 and 67 were sequenced using the Illumina Miseq platform. After low quality sequences and chimeras were removed, the sequence number of each sample was rarefied to 15 131, resulting in 226–579 OTUs at the clustering threshold of 0.97. Our results showed that 99.9% of the sequences belonged to bacteria. SWCNTs addition significantly reduced Shannon diversity and evenness (ANOVA  $P = 0.027$  and  $0.025$ , respectively) on the first day (Table 1), while the richness (OTU and Chao1) did not change ( $P > 0.05$ ). Thereafter, the Shannon indices, OTU richness and evenness were similar among all groups ( $P > 0.05$ ), suggesting a shock loading effect upon SWCNTs addition, followed by a gradual recovery of bacterial diversity. Our results were in accordance with the report by Khodakovskaya et al. that CNTs did not affect soil bacterial diversity after a 9 week influence,<sup>39</sup> whereas Hai et al. reported that continuous addition of 20 mg/L MWCNTs significantly decreased bacterial diversity,<sup>33</sup> indicating the influence of CNTs on microbial diversity was inconclusive.


Pearson correlations between taxa abundances and CNT concentrations were assessed to estimate the impact of SWCNTs on taxonomic groups (Table S2 of the Supporting Information). On Day 1, more taxa were positively affected than negative ones at phylum, class, genus and OTU levels, whereas the opposite trend was observed at all other sampling time. For example, the majority of significantly impacted OTUs (78/106) were of lower abundances on Day 1, yet much smaller fractions of those were reduced on other sampling days (32/155, 27/93, 31/98, 55/126 for Day 20, 40, 60 and 67, respectively). Therefore, the short-term exposure to SWCNTs repressed the growth of the majority of bacterial populations, while continuous interaction exhibited positive effects.

DCA analysis showed that the G1 samples at all sampling time separated from the other groups on a two-axes plot (Figure 3). Hierarchical clustering analysis also showed that the triplicate samples of G1 clustered together and separated from the samples of all other groups (Figure S7 of the Supporting Information). Dissimilarity analyses by Adonis, ANOSIM and MRPP confirmed that G1 was significantly distinct from the other samples (data not shown,  $P < 0.01$ ) at all time points. The changes in bacterial communities, especially the dominant populations, may lead to disturbed system functions.

**Susceptible Genera in Response to SWCNTs.** The influences of SWCNTs on dominant genera (relative abundances over 1%) were further assessed at the initial (Day 1) and final (Day 60) stages of SWCNTs addition. On Day 1, there were 11 dominant genera in all groups, among which only *Zoogloea* exhibited different abundances among groups (ANOVA,  $P < 0.05$ ) and it did not survive at final stage (Figure S8 of the Supporting Information). The  $SV_{30}$  values in our experiment were in the order of G1 (72.8% in average) > G2 (62.4%) > G3  $\approx$  G4  $\approx$  G5 (50.3%) in the first 10 days, which indicated that the addition of SWCNTs improved sludge settling ability (Figure S9 of the Supporting Information). Yin et al. also reported SWCNTs addition improved sludge settling ability by 5.7–10.8% within 5 h.<sup>31</sup> Species of *Zoogloea* (*Zoogloea ramigera*) were widely spread in activated sludge, and have been regarded as the key populations responsible for the flocculation of activated sludges.<sup>37,56,57</sup> Therefore, the

increased *Zoogloea* upon SWCNTs addition in this study might have positively influenced the sludge settling ability.

There were 19 dominant genera on Day 60 (Figure S10 of the Supporting Information), and 7 of them showed significant shifts (Figure 4). *Rudaea*, the most predominant population at



**Figure 4.** Relative abundances of significantly shifted genera in SWCNTs-treated groups compared with control group on Day 60. ANOVA analysis was adopted and the 7 genera had  $P < 0.05$ .

final stage, remarkably increased from 3.17% in G1 to 10.90–26.55% in G2–G5 (Figure 4). The relative abundance of this genus on Day 67 (3.69% in G1, 11.47–26.48% in others) was similar to that of Day 60 (Figure S11 of the Supporting Information). *Rudaea* has been identified in long-term contaminated soils with biphenyl, benzoate and naphthalene, as well as in a petroleum refinery wastewater treatment plant, thus it is of considerable potential for aromatics biodegradation.<sup>58,59</sup> Meanwhile, it is also known for cellulose degradation and was detected as the predominate and sensitive genus susceptible to metal nanoparticles, especially nano-TiO<sub>2</sub>.<sup>60,61</sup> In our study, the increase of *Rudaea* in SWCNTs-treated groups might lead to improved aromatic (such as phenol) degradation capacity.

The abundances of *Burkholderia*, *Singulisphaera*, *Mucilagibacter* and *Labrys* in SWCNTs-treated groups were notably lower than that in G1. Members of *Burkholderia* were known to play important roles in bioremediation of recalcitrant xenobiotics, as well as polyphosphate uptake and accumulation in activated sludge systems.<sup>62–64</sup> Type strains of *Labrys* were reported to be capable of degrading fluorobenzene, chlorobenzene and various pharmaceuticals.<sup>65,66</sup> Thus, the changes of these two bacteria might cause the fluctuation of xenobiotic removal performance. *Singulisphaera* was a newly established genus belonging to the order *Planctomycetales* with biopolymers-degrading ability.<sup>67</sup> Certain species from *Mucilagibacter* including *Mucilagibacter gracilis* and *Mucilagibacter paludis* were also proficient in degrading various biopolymers (pectin, xylan, laminarin, etc.).<sup>68</sup> Therefore, the higher abundances of *Singulisphaera* and *Mucilagibacter* in G1 might result in relatively more EPS degradation, which may explain the higher EPS concentration in SWCNTs-treated groups. *Mobilicoccus* was another increased genus in response to SWCNTs. It has so far only been isolated from fish intestinal tracts and its ecological role in WWTPs kept unknown.<sup>69</sup> *Geothrix* was an anaerobic Fe(III)-reducing bacterium that usually existed in hydrocarbon-contaminated matrix.<sup>70</sup> The roles of this genus also needed further investigation.

*Rhodanobacter* was another predominant genus with relative abundance of over 10%. Although it showed no significant differences among G1 to G4 (ANOVA,  $P > 0.05$ ), it was significantly lower in G5 (7.54%) compared with G1 (16.73%) at the final stage (Figure S11 of the Supporting Information). *Rhodanobacter* was reported to be capable of aromatics degradation and denitrification.<sup>71,72</sup> The decrease in abundance of this genus might negatively affect aromatic degradation and nitrogen removal performance in G5.

Most of the previous studies only searched the shifted microbes upon SWCNTs addition at one time point, yet the long-time and termly detection of community changes could bring us more useful insights. Our results revealed similar responses of *Rudaea*, *Burkholderia*, *Geothrix*, *Mobilicoccus* and *Labrys* at any sampling time, indicating that they were more susceptible to SWCNTs in phenol wastewater treatment systems (Figure S11 of the Supporting Information). However, SWCNTs exhibited varied or converse effects on some other taxa including *Rhodanobacter*, *Singulisphaera* and *Mucilaginibacter* at different time points. It suggested the impacts of SWCNTs were of temporal relations. Under in situ conditions, more complicated surroundings will induce disparate impacts on the community members. Therefore, the ecological effects of SWCNTs should be investigated case by case.

It was previously reported that high concentrations of SWCNTs could significantly reduce urban soil enzyme activity and affect 2,4-dichlorophenol mineralization, possibly by inhibiting the activity of soil endogenous microorganisms, whereas low concentrations of SWCNTs showed no or little influence on the microbes.<sup>19,22</sup> Based on our study, 3.5 g/L seemed to be the threshold for SWCNT toxicity, above which the performance of the SBRs could be unstable upon high phenol influent shock. For controllable engineered system-bioreactors, the performance and stability correlated with functional redundancy.<sup>73–75</sup> Hence, the specific contributions and interactions of the community members, and particularly the impacts of SWCNTs on related functional genes need further investigation to obtain an in-depth understanding of the underlying mechanisms of the SWCNT ecological effects. Because SWCNTs can significantly alter the microbial community structure, both improving and inhibiting wastewater treatment system performance, application of CNTs to wastewater treatment systems should be carefully considered for balancing both positive and negative effects.

**Implications.** Although SWCNTs were shown to be toxic to bacteria elsewhere, this study showed that addition of SWCNTs to phenol wastewater treatment systems could reduce cytotoxicity and increase phenol removal rates. However, the performance of bioreactors receiving a 3.5 g/L dose of SWCNT was significantly lowered upon loading with high concentrations of phenol, suggesting SWCNTs also posed a potential threat to the treatment systems. SWCNTs addition significantly altered the composition and structure of indigenous activated sludge microbial communities. *Zoogloea*, *Rudaea*, *Mobilicoccus*, *Burkholderia*, *Singulisphaera*, *Labrys* and *Mucilaginibacter* were significantly shifted, which might result in microbial community function fluctuations. The high-throughput sequencing technology has proven to be a feasible method for detecting subtle microbial changes resulting from CNT contamination in realistic environmental matrixes, which will contribute to promoting understanding of SWCNT nanotoxicology.

## ■ ASSOCIATED CONTENT

### ■ Supporting Information

MLSS concentration of each group (Figure S1), phenol removal rates on Day 1, 20, 40, 60, 61 and 67 (Figure S2), concentration of bound EPS on Day 3 (Figure S3), fluorescence images of activated sludge on Day 20, 40, 60 and 67 (Figure S4), cell-free DNA fluorescence of on Day 1 and 20 (Figure S5), SEM of G2 (Figure S6), hierarchical clustering analysis of all samples showing (Figure S7), relative abundances of the dominant genera in control and SWCNTs-treated groups on Day 1 and 60 (Figure S8),  $SV_{30}$  value of each group during the first 20 days (Figure S9), relative abundances of the dominant genera in control and SWCNTs-treated groups on Day 60 and 1 (Figure S10), relative abundances of *Rhodanobacter*, *Rudaea*, *Burkholderia*, *Singulisphaera*, *Geothrix*, *Mobilicoccus*, *Labrys* and *Mucilaginibacter* on each sampling time (Figure S11), phenol adsorption by SWCNTs and autoclaved sludge-SWCNTs (Table S1) and number of taxa showing significant correlations with SWCNTs (Table S2). This material is available free of charge via the Internet at <http://pubs.acs.org>.

## ■ AUTHOR INFORMATION

### Corresponding Author

\*Y. Qu. Phone: +86 411 84706251. Fax: +86 411 84706252. E-mail: [qyy@dlut.edu.cn](mailto:qyy@dlut.edu.cn).

### Author Contributions

<sup>||</sup>These authors contributed equally to this article.

### Notes

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

This work is supported by the National Natural Science Foundation of China (No. 21176040), the Program for New Century Excellent Talents in University (No. NCET-13-0077), the Fundamental Research Funds for the Central Universities (No. DUT14YQ107), the Office of the Vice President for Research at the University of Oklahoma and the Collaborative Innovation Center for Regional Environmental Quality. We also acknowledge Dr. Jingwei Wang, Hao Zhou and Zhaojing Zhang (Danlian University of Technology) and Yujia Qin (Oklahoma University) for technical help and useful discussions.

## ■ REFERENCES

- (1) Baughman, R. H.; Zakhidov, A. A.; de Heer, W. A. Carbon nanotubes—The route toward applications. *Science* **2002**, *297* (5582), 787–792.
- (2) Mauter, M. S.; Elimelech, M. Environmental applications of carbon-based nanomaterials. *Environ. Sci. Technol.* **2008**, *42* (16), 5843–5859.
- (3) Petersen, E. J.; Zhang, L. W.; Mattison, N. T.; O'Carroll, D. M.; Whelton, A. J.; Uddin, N.; Nguyen, T.; Huang, Q. G.; Henry, T. B.; Holbrook, R. D.; Chen, K. L. Potential release pathways, environmental fate, and ecological risks of carbon nanotubes. *Environ. Sci. Technol.* **2011**, *45* (23), 9837–9856.
- (4) Neale, P. A.; Jämting, Å. K.; Escher, B. I.; Herrmann, J. A review of the detection, fate and effects of engineered nanomaterials in wastewater treatment plants. *Water Sci. Technol.* **2013**, *68* (7), 1440–1453.
- (5) Sarma, S. J.; Bhattacharya, I.; Brar, S. K.; Tyagi, R. D.; Surampalli, R. Y. Carbon nanotube-bioaccumulation and recent advances in environmental monitoring. *Crit. Rev. Environ. Sci. Technol.* **2015**, *45* (9), 905–938.

- (6) *Nanotoxicology: Materials, Methodologies, and Assessments*; Durán, N., Guterres, S. S., Alves, O. L., Zucolotto, V., Eds.; Springer Science & Business Media: New York, 2013.
- (7) Zhu, B.; Xia, X.; Xia, N.; Zhang, S.; Guo, X. Modification of fatty acids in membranes of bacteria: implication for an adaptive mechanism to the toxicity of carbon nanotubes. *Environ. Sci. Technol.* **2014**, *48*, 4086–4095.
- (8) Kang, S.; Mauter, M. S.; Elimelech, M. Physicochemical determinants of multiwalled carbon nanotube bacterial cytotoxicity. *Environ. Sci. Technol.* **2008**, *42* (19), 7528–7534.
- (9) Kang, S.; Pinault, M.; Pfefferle, L. D.; Elimelech, M. Single-walled carbon nanotubes exhibit strong antimicrobial activity. *Langmuir* **2007**, *23* (17), 8670–8673.
- (10) Pasquini, L. M.; Hashmi, S. M.; Sommer, T. J.; Elimelech, M.; Zimmerman, J. B. Impact of surface functionalization on bacterial cytotoxicity of single-walled carbon nanotubes. *Environ. Sci. Technol.* **2012**, *46* (11), 6297–6305.
- (11) Vecitis, C. D.; Zodrow, K. R.; Kang, S.; Elimelech, M. Electronic-structure-dependent bacterial cytotoxicity of single-walled carbon nanotubes. *ACS Nano* **2010**, *4* (9), 5471–5479.
- (12) Rodrigues, D. F.; Elimelech, M. Toxic effects of single-walled carbon nanotubes in the development of *E. coli* biofilm. *Environ. Sci. Technol.* **2010**, *44* (12), 4583–4589.
- (13) Tu, Y. S.; Lv, M.; Xiu, P.; Huynh, T.; Zhang, M.; Castelli, M.; Liu, Z. R.; Huang, Q.; Fan, C. H.; Fang, H. P.; Zhou, R. H. Destructive extraction of phospholipids from *Escherichia coli* membranes by graphene nanosheets. *Nat. Nanotechnol.* **2013**, *8* (8), 594–601.
- (14) Kang, S.; Mauter, M. S.; Elimelech, M. Microbial cytotoxicity of carbon-based nanomaterials: implications for river water and wastewater effluent. *Environ. Sci. Technol.* **2009**, *43* (7), 2648–2653.
- (15) Tong, Z.; Bischoff, M.; Nies, L. F.; Myer, P.; Applegate, B.; Turco, R. F. Response of soil microorganisms to as-produced and functionalized single-wall carbon nanotubes (SWNTs). *Environ. Sci. Technol.* **2012**, *46* (24), 13471–13479.
- (16) Farré, M.; Gajda-Schrantz, K.; Kantiani, L.; Barceló, D. Ecotoxicity and analysis of nanomaterials in the aquatic environment. *Anal. Bioanal. Chem.* **2009**, *393* (1), 81–95.
- (17) Eckelman, M. J.; Mauter, M. S.; Isaacs, J. A.; Elimelech, M. New perspectives on nanomaterial aquatic ecotoxicity: Production impacts exceed direct exposure impacts for carbon nanotubes. *Environ. Sci. Technol.* **2012**, *46* (5), 2902–2910.
- (18) Rodrigues, D. F.; Jaisi, D. P.; Elimelech, M. Toxicity of functionalized single-walled carbon nanotubes on soil microbial communities: Implications for nutrient cycling in soil. *Environ. Sci. Technol.* **2012**, *47* (1), 625–633.
- (19) Jin, L.; Son, Y.; Yoon, T. K.; Kang, Y. J.; Kim, W.; Chung, H. High concentrations of single-walled carbon nanotubes lower soil enzyme activity and microbial biomass. *Ecotoxicol. Environ. Saf.* **2013**, *88* (1), 9–15.
- (20) Chung, H.; Son, Y.; Yoon, T. K.; Kim, S.; Kim, W. The effect of multi-walled carbon nanotubes on soil microbial activity. *Ecotoxicol. Environ. Saf.* **2011**, *74* (4), 569–575.
- (21) Shrestha, B.; Acosta-Martinez, V.; Cox, S. B.; Green, M. J.; Li, S.; Cañas-Carrell, J. E. An evaluation of the impact of multiwalled carbon nanotubes on soil microbial community structure and functioning. *J. Hazard. Mater.* **2013**, *261*, 188–197.
- (22) Zhou, W. Q.; Shan, J.; Jiang, B. Q.; Wang, L. H.; Feng, J. F.; Guo, H. Y.; Ji, R. Inhibitory effects of carbon nanotubes on the degradation of <sup>14</sup>C-2,4-dichlorophenol in soil. *Chemosphere* **2013**, *90* (2), 527–534.
- (23) Cui, X. Y.; Jia, F.; Chen, Y. X.; Gan, J. Influence of single-walled carbon nanotubes on microbial availability of phenanthrene in sediment. *Ecotoxicology* **2011**, *20* (6), 1277–1285.
- (24) Musee, N.; Thwala, M.; Nota, N. The antibacterial effects of engineered nanomaterials: implications for wastewater treatment plants. *J. Environ. Monit.* **2011**, *13* (5), 1164–1183.
- (25) Upadhyayula, V. K. K.; Deng, S.; Mitchell, M. C.; Smith, G. B. Application of carbon nanotube technology for removal of contaminants in drinking water: A review. *Sci. Total Environ.* **2009**, *408* (1), 1–13.
- (26) Brar, S. K.; Verma, M.; Tyagi, R. D.; Surampalli, R. Y. Engineered nanoparticles in wastewater and wastewater sludge—Evidence and impacts. *Waste Manage.* **2010**, *30* (3), 504–520.
- (27) Chen, Y. G.; Wang, D. B.; Zhu, X. Y.; Zheng, X.; Feng, L. Y. Long-term effects of copper nanoparticles on wastewater biological nutrient removal and N<sub>2</sub>O generation in the activated sludge process. *Environ. Sci. Technol.* **2012**, *46* (22), 12452–12458.
- (28) Yang, Y.; Quensen, J.; Mathieu, J.; Wang, Q.; Wang, J.; Li, M. Y.; Tiedje, J. M.; Alvarez, P. J. J. Pyrosequencing reveals higher impact of silver nanoparticles than Ag<sup>+</sup> on the microbial community structure of activated sludge. *Water Res.* **2014**, *48*, 317–325.
- (29) Goyal, D.; Zhang, X. J.; Rooney-Varga, J. N. Impacts of single-walled carbon nanotubes on microbial community structure in activated sludge. *Letts. Appl. Microbiol.* **2010**, *51* (4), 428–435.
- (30) Luongo, L. A.; Zhang, X. J. Toxicity of carbon nanotubes to the activated sludge process. *J. Hazard. Mater.* **2010**, *178* (1–3), 356–362.
- (31) Yin, Y.; Zhang, X. J.; Graham, J.; Luongo, L. Examination of purified single-walled carbon nanotubes on activated sludge process using batch reactors. *J. Environ. Sci. Health, Part A: Toxic/Hazard. Subst. Environ. Eng.* **2009**, *44* (7), 661–665.
- (32) Yin, Y.; Zhang, X. J. Evaluation of the impact of single-walled carbon nanotubes in an activated sludge wastewater reactor. *Water Sci. Technol.* **2008**, *58* (3), 623–628.
- (33) Hai, R.; Wang, Y.; Wang, X.; Du, Z.; Li, Y. Impacts of multiwalled carbon nanotubes on nutrient removal from wastewater and bacterial community structure in activated sludge. *PLoS One* **2014**, *9* (9), e107345.
- (34) Li, L. L.; Tong, Z. H.; Fang, C. Y.; Chu, J.; Yu, H. Q. Response of anaerobic granular sludge to single-wall carbon nanotube exposure. *Water Res.* **2015**, *70*, 1–8.
- (35) Jin, L.; Son, Y.; DeForest, J. L.; Kang, Y. J.; Kim, W.; Chung, H. Single-walled carbon nanotubes alter soil microbial community composition. *Sci. Total Environ.* **2014**, *466*, 533–538.
- (36) Luo, C.; Tsementzi, D.; Kyrpidis, N.; Read, T.; Konstantinidis, K. T. Direct comparisons of Illumina vs. Roche 454 sequencing technologies on the same microbial community DNA sample. *PLoS One* **2012**, *7* (2), e30087.
- (37) Zhang, T.; Shao, M. F.; Ye, L. 454 Pyrosequencing reveals bacterial diversity of activated sludge from 14 sewage treatment plants. *ISME J.* **2012**, *6* (6), 1137–1147.
- (38) Ge, Y.; Schimel, J. P.; Holden, P. A. Identification of soil bacteria susceptible to TiO<sub>2</sub> and ZnO nanoparticles. *Appl. Environ. Microbiol.* **2012**, *78* (18), 6749–6758.
- (39) Khodakovskaya, M. V.; Kim, B. S.; Kim, J. N.; Alimohammadi, M.; Dervishi, E.; Mustafa, T.; Cernigla, C. E. Carbon nanotubes as plant growth regulators: Effects on tomato growth, reproductive system, and soil microbial community. *Small* **2013**, *9* (1), 115–123.
- (40) Shen, E.; Qu, Y. Y.; Zhou, H.; Kong, C. L.; Ma, Q.; Zhang, X. W.; Zhou, J. T. Catalytic performance and stability of C-C bond hydrolase BphD immobilized onto single-wall carbon nanotubes. *Chin. J. Catal.* **2013**, *34* (4), 723–733.
- (41) Raunkjær, K.; Hvitved-Jacobsen, T.; Nielsen, P. H. Measurement of pools of protein, carbohydrate and lipid in domestic wastewater. *Water Res.* **1994**, *28* (2), 251–262.
- (42) Purkhold, U.; Pommerening-Röser, A.; Juretschko, S.; Schmid, M. C.; Koops, H. P.; Wagner, M. Phylogeny of all recognized species of ammonia oxidizers based on comparative 16S rRNA and *amoA* sequence analysis: Implications for molecular diversity surveys. *Appl. Environ. Microbiol.* **2000**, *66*, 5368–5382.
- (43) Ma, Q.; Qu, Y.; Shen, W.; Zhang, Z.; Wang, J.; Liu, Z.; Li, D.; Li, H.; Zhou, J. Bacterial community compositions of coking wastewater treatment plants in steel industry revealed by Illumina high-throughput sequencing. *Bioresour. Technol.* **2015**, *179*, 436–443.
- (44) Bates, S. T.; Berg-Lyons, D.; Caporaso, J. G.; Walters, W. A.; Knight, R.; Fierer, N. Examining the global distribution of dominant archaeal populations in soil. *ISME J.* **2010**, *5* (5), 908–917.

- (45) Magoč, T.; Salzberg, S. L. FLASH: Fast length adjustment of short reads to improve genome assemblies. *Bioinformatics* **2011**, *27*, 2957–2963.
- (46) Deng, Y.; He, Z.; Xu, M.; Qin, Y.; Van Nostrand, J. D.; Wu, L.; Roe, B. A.; Wiley, G.; Hobbie, S. E.; Reich, P. B.; Zhou, J. Elevated carbon dioxide alters the structure of soil microbial communities. *Appl. Environ. Microbiol.* **2012**, *78* (8), 2991–2995.
- (47) Edgar, R. C.; Haas, B. J.; Clemente, J. C.; Quince, C.; Knight, R. UCHIME improves sensitivity and speed of chimera detection. *Bioinformatics* **2011**, *27* (16), 2194–2200.
- (48) Li, W. Z.; Godzik, A. CD-HIT: A fast program for clustering and comparing large sets of protein or nucleotide sequences. *Bioinformatics* **2006**, *22* (13), 1658–1659.
- (49) Wang, Q.; Garrity, G. M.; Tiedje, J. M.; Cole, J. R. Naïve Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Appl. Environ. Microbiol.* **2007**, *73* (16), 5261–5267.
- (50) Bour, A.; Mouchet, F.; Silvestre, J.; Gauthier, L.; Pinelli, E. Environmentally relevant approaches to assess nanoparticles ecotoxicity: A review. *J. Hazard. Mater.* **2015**, *283*, 764–777.
- (51) Nowack, B.; David, R. M.; Fissan, H.; Morris, H.; Shatkin, J. A.; Stintz, M.; Zepp, R.; Brouwer, D. Potential release scenarios for carbon nanotubes used in composites. *Environ. Int.* **2013**, *59*, 1–11.
- (52) Ren, X. M.; Chen, C. L.; Nagatsu, M.; Wang, X. K. Carbon nanotubes as adsorbents in environmental pollution management: A review. *Chem. Eng. J.* **2011**, *170* (2–3), 395–410.
- (53) Parks, A. N.; Chandler, G. T.; Portis, L. M.; Sullivan, J. C.; Perron, M. M.; Cantwell, M. G.; Burgess, R. M.; Ho, K. T.; Ferguson, P. L. Effects of single-walled carbon nanotubes on the bioavailability of PCBs in field-contaminated sediments. *Nanotoxicology* **2013**, *8* (Suppl 1), 111–117.
- (54) Lin, D.; Xingt, B. Adsorption of phenolic compounds by carbon nanotubes: Role of aromaticity and substitution of hydroxyl groups. *Environ. Sci. Technol.* **2008**, *42* (19), 7254–7259.
- (55) Qu, X.; Alvarez, P. J.; Li, Q. Applications of nanotechnology in water and wastewater treatment. *Water Res.* **2013**, *47* (12), 3931–3946.
- (56) Rosselló-Mora, R. A.; Wagner, M.; Amann, R.; Schleifer, K. H. The abundance of *Zoogloea ramigera* in sewage treatment plants. *Appl. Environ. Microbiol.* **1995**, *61* (2), 702–707.
- (57) Salehizadeh, H.; Shojaosadati, S. A. Extracellular biopolymeric flocculants: Recent trends and biotechnological importance. *Biotechnol. Adv.* **2001**, *19* (5), 371–385.
- (58) Uhlik, O.; Wald, J.; Strejcek, M.; Musilova, L.; Ridl, J.; Hroudova, M.; Vlcek, C.; Cardenas, E.; Mackova, M.; Macek, T. Identification of bacteria utilizing biphenyl, benzoate, and naphthalene in long-term contaminated soil. *PLoS One* **2012**, *7* (7), e40653.
- (59) Ibarbalz, F. M.; Figuerola, E. L.; Erijman, L. Industrial activated sludge exhibit unique bacterial community composition at high taxonomic ranks. *Water Res.* **2013**, *47* (11), 3854–3864.
- (60) Weon, H. Y.; Yoo, S. H.; Kim, Y. J.; Lee, C. M.; Kim, B. Y.; Jeon, Y. A.; Hong, S. B.; Anandham, R.; Kwon, S. W. *Rudaea cellulositytica* gen. nov., sp. nov., isolated from soil. *Int. J. Syst. Evol. Microbiol.* **2009**, *59*, 2308–2312.
- (61) Shah, V.; Jones, J.; Dickman, J.; Greenman, S. Response of soil bacterial community to metal nanoparticles in biosolids. *J. Hazard. Mater.* **2014**, *274*, 399–403.
- (62) Coenye, T.; Vandamme, P. Diversity and significance of *Burkholderia* species occupying diverse ecological niches. *Environ. Microbiol.* **2003**, *5* (9), 719–729.
- (63) Wang, J.; Quan, X.; Wu, L.; Qian, Y.; Werner, H. Bioaugmentation as a tool to enhance the removal of refractory compound in coke plant wastewater. *Process Biochem.* **2002**, *38* (5), 777–781.
- (64) Mullan, A.; Quinn, J. P.; McGrath, J. W. Enhanced phosphate uptake and polyphosphate accumulation in *Burkholderia cepacia* grown under low-pH conditions. *Microb. Ecol.* **2002**, *44* (1), 69–77.
- (65) Carvalho, M. F.; De Marco, P.; Duque, A. F.; Pacheco, C. C.; Janssen, D. B.; Castro, P. M. *Labrys portucalensis* sp. nov., a fluorobenzene-degrading bacterium isolated from an industrially contaminated sediment in northern Portugal. *Int. J. Syst. Evol. Microbiol.* **2008**, *58* (3), 692–698.
- (66) Amorim, C. L.; Moreira, I. S.; Maia, A. S.; Tiritan, M. E.; Castro, P. M. Biodegradation of ofloxacin, norfloxacin, and ciprofloxacin as single and mixed substrates by *Labrys portucalensis* F11. *Appl. Microbiol. Biotechnol.* **2014**, *98* (7), 3181–3190.
- (67) Kulichevskaya, I. S.; Ivanova, A. O.; Baulina, O. I.; Bodelier, P. L.; Damsté, J. S. S.; Dedysh, S. N. *Singulisphaera acidiphila* gen. nov., sp. nov., a non-filamentous, Isosphaera-like planctomycete from acidic northern wetlands. *Int. J. Syst. Evol. Microbiol.* **2008**, *58* (5), 1186–1193.
- (68) Pankratov, T. A.; Tindall, B. J.; Liesack, W.; Dedysh, S. N. *Mucilaginibacter paludis* gen. nov., sp. nov. and *Mucilaginibacter gracilis* sp. nov., pectin-, xylan- and laminarin-degrading members of the family *Sphingobacteriaceae* from acidic Sphagnum peat bog. *Int. J. Syst. Evol. Microbiol.* **2007**, *57* (10), 2349–2354.
- (69) *The Prokaryotes: Actinobacteria*, 4th, ed.; Rosenberg, E., DeLong, E. F., Lory, S., Stackebrandt, E., Thompson, F., Eds.; Springer: Berlin/Heidelberg, 2014.
- (70) Coates, J. D.; Ellis, D. J.; Gaw, C. V.; Lovley, D. R. *Geothrix fermentans* gen. nov., sp. nov., a novel Fe (III)-reducing bacterium from a hydrocarbon-contaminated aquifer. *Int. J. Syst. Evol. Microbiol.* **1999**, *49* (4), 1615–1622.
- (71) Felföldi, T.; Székely, A. J.; Gorál, R.; Barkács, K.; Scheirich, G.; András, J.; Rácz, A.; Márialigeti, K. Polyphasic bacterial community analysis of an aerobic activated sludge removing phenols and thiocyanate from coke plant effluent. *Bioresour. Technol.* **2010**, *101* (10), 3406–3414.
- (72) Bacosa, H. P.; Suto, K.; Inoue, C. Bacterial community dynamics during the preferential degradation of aromatic hydrocarbons by a microbial consortium. *Int. Biodeterior. Biodegrad.* **2012**, *74*, 109–115.
- (73) Briones, A.; Raskin, L. Diversity and dynamics of microbial communities in engineered environments and their implications for process stability. *Curr. Opin. Biotechnol.* **2003**, *14* (3), 270–276.
- (74) Pholchan, M. K.; Baptista, J. de C.; Davenport, R. J.; Curtis, T. P. Systematic study of the effect of operating variables on reactor performance and microbial diversity in laboratory-scale activated sludge reactors. *Water Res.* **2010**, *44* (5), 1341–1352.
- (75) Cabrol, L.; Malhautier, L.; Poly, F.; Lepeuple, A. S.; Fanlo, J. L. Bacterial dynamics in steady-state biofilters: Beyond functional stability. *FEMS Microbiol. Ecol.* **2012**, *79* (1), 260–271.