

Response of *Desulfovibrio vulgaris* to Alkaline Stress^{∇†}

Sergey Stolyar,^{1,10*} Qiang He,^{6,10}§ Marcin P. Joachimiak,^{2,10} Zhili He,¹⁰ Zamin Koo Yang,^{4,10}
Sharon E. Borglin,^{9,10} Dominique C. Joyner,^{9,10} Katherine Huang,^{2,10} Eric Alm,^{7,10}
Terry C. Hazen,^{9,10} Jizhong Zhou,^{4,8,10} Judy D. Wall,^{5,10} Adam P. Arkin,^{2,3,10}
and David A. Stahl^{1,10}

Department of Civil and Environmental Engineering, University of Washington, Seattle, Washington¹; Physical Bioscience Division, Lawrence Berkeley National Laboratory, Berkeley, California²; Department of Bioengineering, University of California, Berkeley, California³; Oak Ridge National Laboratory, Oak Ridge, Tennessee⁴; Department of Biochemistry, University of Missouri-Columbia, Columbia, Missouri⁵; Department of Civil and Environmental Engineering, Temple University, Philadelphia, Pennsylvania⁶; Department of Civil and Environmental Engineering, Massachusetts Institute of Technology, Cambridge, Massachusetts⁷; Institute for Environmental Genomics, Department of Botany and Microbiology, University of Oklahoma, Norman, Oklahoma⁸; Earth Science Division, Lawrence Berkeley National Laboratory, Berkeley, California⁹; and Virtual Institute for Microbial Stress and Survival¹⁰‡

Received 21 February 2007/Accepted 10 September 2007

The response of exponentially growing *Desulfovibrio vulgaris* Hildenborough to pH 10 stress was studied using oligonucleotide microarrays and a study set of mutants with genes suggested by microarray data to be involved in the alkaline stress response deleted. The data showed that the response of *D. vulgaris* to increased pH is generally similar to that of *Escherichia coli* but is apparently controlled by unique regulatory circuits since the alternative sigma factors (sigma S and sigma E) contributing to this stress response in *E. coli* appear to be absent in *D. vulgaris*. Genes previously reported to be up-regulated in *E. coli* were up-regulated in *D. vulgaris*; these genes included three ATPase genes and a tryptophan synthase gene. Transcription of chaperone and protease genes (encoding ATP-dependent Clp and La proteases and DnaK) was also elevated in *D. vulgaris*. As in *E. coli*, genes involved in flagellum synthesis were down-regulated. The transcriptional data also identified regulators, distinct from sigma S and sigma E, that are likely part of a *D. vulgaris* Hildenborough-specific stress response system. Characterization of a study set of mutants with genes implicated in alkaline stress response deleted confirmed that there was protective involvement of the sodium/proton antiporter NhaC-2, tryptophanase A, and two putative regulators/histidine kinases (DVU0331 and DVU2580).

Sulfate-reducing bacteria (SRB) are ubiquitous in nature and play an important role in global carbon and sulfur cycling. Their habitat range includes freshwater, marine, and hypersaline aquatic systems, cold oceanic sediments, the deep subsurface, hydrothermal vents, and hot springs (11, 26, 35). Although long thought to have a relatively restricted catabolic range, this functionally defined assemblage is now recognized to be remarkably versatile. SRB mediate the degradation of aromatic compounds once thought to be refractory to anaerobic degradation, including benzene (2, 3, 8, 22, 23), and reduce a variety of metals, including radionuclides (19, 21, 24, 35). For these reasons they have also been studied for possible use in the bioremediation of environments contaminated with organic and metal pollutants.

Desulfovibrio vulgaris Hildenborough is one of the better-characterized SRB. This gram-negative deltaproteobacterium, isolated 60 years ago from clay soil in Hildenborough, Kent

(United Kingdom), has served as one of the principal models for resolving the physiological and genetic basis of sulfate respiration. The recent completion of its genome sequence (14) has enabled genome-wide expression studies (6, 13, 27, 36, 37) that are now beginning to resolve its adaptive response to changing environmental parameters. Although this information is essential for predicting its behavior in possible applications for bioremediation, information about the range of conditions that support *D. vulgaris* growth or survival remains scarce.

Alkaline environments are common in nature (e.g., alkaline groundwater, lakes, and intestinal segments of some higher organisms) and in sites contaminated by human activity (29). There are some data documenting the presence of SRB in alkaline environments (1), but little is known about specific adaptive mechanisms of these bacteria or even whether mechanisms common to better-characterized organisms, such as *Escherichia coli* and *Bacillus subtilis*, are used by SRB. The adaptive strategies used by other microbes include (i) increased proton pumping by ATP synthase, (ii) increased metabolic acid production through amino acid deaminases and sugar fermentation, (iii) changes in cell surface properties, and (iv) increased expression and activity of monovalent cation/proton antiporters (9, 29, 33). Among these strategies, monovalent cation/proton antiporters are thought to play a central role in alkaline pH homeostasis in many bacteria (29).

In order to better resolve similarities and differences in the

* Corresponding author. Mailing address: University of Washington, Department of Civil and Environmental Engineering, 616 NE North Lake Place, Box 355014, Seattle, WA 98195-2700. Phone: (206) 543-2094. Fax: (206) 685-9185. E-mail: sstolyar@u.washington.edu.

† Supplemental material for this article may be found at <http://jb.asm.org/>.

§ Present address: Department of Civil and Environmental Engineering, University of Tennessee, Knoxville, TN.

‡ <http://vimss.lbl.gov>.

∇ Published ahead of print on 5 October 2007.

TABLE 1. Strains of *D. vulgaris* Hildenborough used in this study

Strain	Putative gene	Annotation	Strain description	Source
ATCC 29579	Not applicable	Wild type	Parent of deletion and plasmid insertion mutants	ATCC
JW381	DVU3108	<i>nhaC</i>	$\Delta nhaC::ntp$ MP ⁺	This study
JW391	DVU2204	<i>tnaA</i>	<i>tnaA::pMO391</i> MP ⁺ ^b	This study
JW3024	DVU2580	Regulator	Tn5-RL27 insertion ^c	This study
JW3011	DVU0331	Histidine kinase	Tn5-RL27 insertion	This study
JW801	Megaplasmid deleted (MP ⁻) ^a	Wild type	Parent strain of transposon mutants	G. Voordow

^a MP, 202-kb megaplasmid endogenous to *D. vulgaris* Hildenborough (= ATCC 29579).

^b pMO391 is pCR8/GW/TOPO (Invitrogen) with an internal *tnaA* fragment from *D. vulgaris* used to create an insertional mutation through homologous recombination (Sp⁺).

^c Tn5-RL27 is a modified high-frequency insertion transposon (19).

adaptation of *Desulfovibrio* species, we used genome-wide transcription profiling to characterize the response of *D. vulgaris* to an upshift in the pH of its growth medium. Adaptive mechanisms suggested by transcriptional analysis were then examined by characterizing a study set of mutants with genes implicated in the alkaline stress response deleted. Together, these analyses revealed a response system mechanistically similar to the systems of better-characterized species.

MATERIALS AND METHODS

Cell growth and pH upshift conditions. Two-liter bottles containing 1.8 liters of defined piperazine-*N,N'*-bis(2-ethanesulfonic acid) (PIPES)-buffered medium LS4D (13, 27) amended with 60 mM lactate and 50 mM sulfate were inoculated with exponentially grown cultures (1:10) to generate three biological replicate pairs consisting of a control and a test culture. The headspace of the vessels was 5% H₂, 5% CO₂, and 90% N₂. An alkaline stress (pH 10) was applied to each test culture at mid-exponential growth (optical density at 600 nm, ca. 0.4) by addition of 35 ml of 2 M KOH with vigorous stirring. The control of each pair was treated by addition of 35 ml of sterile anoxic distilled water. Samples (300 ml) were taken for the gene expression analysis from each bottle at 30, 60, 120, and 240 min after addition of the stressor. The zero time samples were taken prior to stressor addition. The sample biomass was immediately chilled to 4°C during collection by pumping the sample out of the vessel with a peristaltic pump through a 7-m-long stainless steel tube immersed in an ice bath. Following centrifugation at 11,000 × *g* for 10 min at 4°C and decanting of the medium, the cell pellet was resuspended in 50 ml of anoxic 10 mM phosphate-buffered saline and cells were collected again by centrifugation. The cell pellets were immediately frozen in liquid N₂ and stored at -80°C until RNA isolation.

To determine the growth rates of mutants at different pH values, cells were grown in B3 medium (containing [per liter] 0.1 g of NaCl, 0.1 g of MgCl₂ · 6H₂O, 0.1 g of CaCl₂ · 2H₂O, 0.5 g of NH₄Cl, 0.1 g of KCl, 1.4 g of Na₂SO₄, 1 g of Na₂S, 0.001 g of resazurin, 1 ml of 1 M K₂HPO₄, 1 ml of trace minerals, 1 ml of Thauer's vitamins, 1 ml of 1 M cysteine, and 1 ml of 1 M Na₂S). The following buffering agents were used: 25 mM sodium bicarbonate for pH 7; 50 mM Tris for pH 7.5; and 50 mM glycine for pH 8.0 and 9.0. In all media 50 mM lactate and 40 mM sodium sulfate were used.

Analytical methods. The concentrations of organic acids (lactate, pyruvate, acetate, formate, and fumarate) and inorganic ions (sulfate and phosphate) in culture media were determined using a Dionex 500 system equipped with an AS11HC column. In some cases the concentrations of organic acids were also measured using a high-performance liquid chromatograph equipped with an HPX 78 (Bio-Rad) column. Hydrogen concentrations were determined with an RGD2 reduction gas detector (Trace Analytical) with a 60/80 Mole Sieve 5A column (6 ft by 0.125 in.) with N₂ as the carrier gas.

Mutant construction. Bacterial mutants and strains are listed in Table 1. Deletion mutant JW381 ($\Delta nhaC-2$) was constructed by marker exchange with a mutagenic plasmid as described elsewhere (4). The primers used for mutagenic plasmid construction were 5'-TATGGCAGATGTCAATGCCGAAGT-3' and 5'-AAGACTGTAGCCGTACCTCGAATCTAATGTAGGCTCCAGTGGCCGA-3' for the upstream region and 5'-ACGGCTTCCACGTCAACTATCTCA-3' and 5'-AATCCGCTCACTAAGTTCATAGACCGTAGGGAAGGGCTA CCTGAGGC-3' for the DNA region downstream of the gene. The kanamycin resistance marker replacing *nhaC-2* is also flanked by sequence bar codes unique

to this deletion, with 5'-GCCGACAGAGCTTGAGATA-3' at the promoter-proximal end of the kanamycin marker and 5'-AGCCTGGAACAGCTATACA C-3' at the distal end.

Insertion mutant JW391 was constructed using plasmid pMO391 bearing a spectinomycin resistance gene (Sp^r). pMO391 is pCR8/GW/TOPO (Invitrogen) with an internal *tnaA* (DVU2204) fragment from *D. vulgaris* that was used to create an insertional mutation of *tnaA* through homologous recombination. The primers used for strain construction were tnaA350Fd (5'-ACAAGCCCGTCTT CATCTCCAAC-3') (forward) and tnaA1273Rv (5'-TGTAGTCCATGTGGT CGTTGGTGT-3') (reverse). The transposon mutants were generated by conjugation between *D. vulgaris* and *E. coli* BW20767(pRL27) (19). The conjugation procedure was a modification of the method of Fu and Voordouw (12). Cultures of *D. vulgaris* were grown to mid-exponential phase and combined at a 3:1 or 6:1 ratio with the *E. coli* donor grown to early exponential phase in LC medium (1.0% [wt/vol] tryptone, 0.5% [wt/vol] yeast extract, 0.5% [wt/vol] NaCl). Mating mixtures were concentrated by centrifugation. The concentrated cells were placed onto filter disks (pore size, 0.22 μm; GSWP; Millipore Billerica, MA), and the disks were placed on the surface of solidified LS4 (LS4D with 1% [wt/vol] yeast extract added) and incubated for 16 h at 34°C. Then the cells were washed from the membrane with 2 ml LS4 medium. After 6 h of incubation, antibiotic G418 (400 μg/ml) was added to select for the transposon mutants and nalidixic acid (200 μg/ml) was added to select against the *E. coli* donor. Then cells were spread onto LS4 agar (100 to 500 μl/plate) with both antibiotics. The plates were incubated in the anaerobic chamber at 34°C for at least 4 days to allow the colonies to grow. The chromosomal localization of the transposon insertions was identified by sequencing DNA after semirandom PCR amplification. For semirandom PCR, a variation of a protocol described by Chun et al. (7) was used. One microliter of a 50-μl boiled single-colony suspension in distilled H₂O was used as the template DNA in a 20-μl PCR mixture containing primer tprRL17-1 (5'-AACAAGCCAGGGATGTAAACG-3') (19) and either primer CEKG 2A (5'-GGCCACGCGTTCGACTAGTACN10AGAG-3'), primer CEKG 2B (5'-GGCCACGCGTTCGACTAGTACN10ACGCC-3'), or primer CEKG 2C (5'-GGCCACGCGTTCGACTAGTACN10GATAT-3'). One microliter of a 1:5 dilution of this reaction mixture was used as the template DNA for a second PCR performed with primers tprRL17-2 (5'-AGCCCTTAGGCCTCTCAAAGCAA-3') and CEKG 4 (5'-GGCCACGCGTTCGACTAGTAC-3'). For the first reaction, the thermocycler conditions were 94°C for 2 min, followed by six cycles of 94°C for 30 s, 42°C for 30 s (with the temperature reduced 1°C per cycle), and 72°C for 3 min and then 25 cycles of 94°C for 30 s, 65°C for 30 s, and 72°C for 3 min; for the second reaction, the thermocycler conditions were 30 cycles of 94°C for 30 s, 65°C for 30 s, and 72°C for 3 min. Samples that produced distinct bands on an agarose gel after the second reaction were cleaned with a PCR purification kit (Qiagen, Valencia, CA) and sequenced by using primer tprRL17-1. The chromosomal locations of the insertions were identified by BLAST analysis of the sequences adjacent to the transposon compared with the complete genome.

Microarray construction, hybridization, and image analysis. DNA microarrays covering 3,482 of the 3,551 protein-encoding sequences of the *D. vulgaris* genome annotated by TIGR (14) were designed, constructed, and validated with 70-mer oligonucleotide probes as previously described (6, 13, 27). Briefly, all oligonucleotides were commercially synthesized without modification by MWG Biotech Inc. (High Point, NC), prepared in 50% (vol/vol) dimethyl sulfoxide (Sigma-Aldrich, St. Louis, MO), and spotted onto UltraGAPS glass slides (Corning Life Sciences, Corning, NY) using a BioRobotics Microgrid II microarrayer (Genomic Solutions, Ann Arbor, MI). For each oligonucleotide probe there

were two replicates on a single slide. After printing, the oligonucleotide probes were fixed onto the slides by UV cross-linking (600 mJ of energy) according to the protocol of the manufacturer of the UltraGAPS glass slides (Corning Life Science).

Total RNA extraction, purification, and labeling were performed independently for each cell sample using previously described protocols (6, 13). Briefly, total cellular RNA was isolated using TRIzol reagent (Invitrogen, Carlsbad, CA) by following the manufacturer's protocol. RNA extracts were purified according to the RNeasy mini kit (Qiagen, Valencia, CA) instructions, and on-column DNase digestion was performed with the RNase-free DNase Set (Qiagen) to remove genomic DNA contamination according to the manufacturer's procedure. Labeling of cDNA targets from purified total RNA was carried out using the reverse transcriptase reaction with random hexamer priming and the fluorophore Cy5-dUTP (Amersham Biosciences, Piscataway, NJ). Genomic DNA was extracted from *D. vulgaris* cultures at stationary phase and labeled with the fluorophore Cy3-dUTP (Amersham Biosciences). The efficiency of labeling was routinely monitored by measuring the absorbance at 260 nm (for DNA concentration), 550 nm (for Cy3), or 650 nm (for Cy5).

To hybridize a single glass slide, the Cy5-dUTP-labeled cDNA probes obtained from stressed or unstressed cultures were mixed with equal amounts of the Cy3-dUTP-labeled genomic DNA (6, 27). Labeled genomic DNA (Cy3) was used as a control and as the common reference to cohybridize with labeled RNA (Cy5) samples for each slide. After washing and drying, the microarray slides were scanned using the ScanArray Express microarray analysis system (Perkin Elmer). The fluorescence intensity of both the Cy5 and Cy3 fluorophores was analyzed with the software ImaGene, version 6.0 (Biodiscovery, Marina Del Rey, CA).

Microarray analysis. Arrays were scanned using the scanning laser confocal fluorescence microscope of the ScanArray microarray analysis system (GSI Lumonics), and hybridization signal intensities were quantitated using the software ImaGene (Biodiscovery). Statistical analysis of the microarray data was performed as described elsewhere (6), and cluster analysis was performed using TIGR Multi Experiment Viewer (MeV) (32). To assess the significance of the normalized log ratio, a Z score was calculated by using the following equation:

$$Z = \frac{\log_2(\text{treatment/control})}{\sqrt{0.25 + \Sigma\text{variance}}}$$

where 0.25 is a pseudovariance term. Log ratios and Z values for all genes detected in this study are listed in Table S1 in the supplemental material. For the microarray experiments, each \log_2 value represents the average of the results of three hybridization experiments using cDNA derived from two different cultures of *D. vulgaris*.

Protein sequence analysis. For protein sequence analysis, the programs Tmpred (http://www.ch.embnet.org/software/TMPRED_form.html) (15), LipoP 1.0 Server (<http://www.cbs.dtu.dk/services/LipoP>) (17), and SignalP 3.0 (10) from the ExPASy database were used. The genomes and open reading frames of *D. vulgaris* and the gene contents of different sequenced genomes were analyzed using Microbesonline (<http://www.microbesonline.org/>) and NCBI databases.

Microarray data accession number. All microarray data have been deposited in the NCBI Gene Expression Omnibus (GEO) database under accession number GSE8197.

RESULTS AND DISCUSSION

General features of the physiological and transcriptional response to elevated pH. The initial studies examined the general physiological response of *D. vulgaris* to an abrupt increase in pH (Fig. 1). Lactate consumption stopped immediately following the pH shift, as did production of acetate (Fig. 1B). Hydrogen accumulation was not detected during the stress. However, viable cell numbers remained constant during the 2 h following the shift (Fig. 1A), as demonstrated by enumerating cells on agar plates and immediate resumption of growth measured by optical density following transfer to fresh neutral medium. During this period of exposure to elevated pH, altered expression of approximately 400 genes was observed. At 30, 60, 120, and 240 min following the shift, transcription of 78, 122, 178, and 184 genes, respectively, was significantly up-




FIG. 1. Growth and pH changes during pH upshift. (A) Cell numbers. (B) Lactate and acetate concentrations in the growth medium of the stressed culture. The data are means of triplicate cultures, and the observed variability was within 10%. The error bars indicate standard deviations of the means ($n = 3$).

regulated (\log_2 signal intensity change greater than 1 and Z score greater than 2). The transcription of comparable numbers of genes was down-regulated at least twofold at the same time points (175, 267, 210, and 183 genes, respectively). The majority of the up-regulated genes fell into the following COG functional categories: amino acid biosynthesis, energy metabolism, and signal transduction systems. A large fraction of the down-regulated genes were assigned to signal transduction, transcription, and phage-related categories.

Using K-means clustering analysis with the Euclidian distance, all genes were assigned to 30 groups based on their patterns of expression. Graphs representing all groups are shown in Fig. S1 in the supplemental material. Genes comprising each group are listed in Table S2 in the supplemental material. In our initial analysis we focused on cluster 22 (44 genes), which was comprised of genes that were the most highly up-regulated at all time points; cluster 19, which consisted of genes that were only moderately up-regulated (122 genes); cluster 4, which was comprised of the most highly down-regulated genes (57 genes); and cluster 21, which consisted of genes which were moderately down-regulated (100 genes) at all time points.

Genes up-regulated during exposure to high pH. Elevated pH is generally thought to stress the cell through alkalinization of the cytoplasm, reduction of the membrane potential, and damage to proteins and the cell envelope (9, 33; for reviews, see references 29 and 31). The cell responds by pumping protons inside, importing or synthesizing compounds to acidify the cytoplasm, and activating systems of protein repair or degradation. These processes are all reflected in the response of *E. coli* to elevated pH. In alkali-stressed *D. vulgaris*, genes encoding ATPase synthase, Na^+/H^+ antiporter NhaC-2 (DVU3108), chaperones and proteases such as DnaK (DVU0811; $\log_2 R = 0.8$ to 1.5), ATP-dependent Clp protease subunit B (DVU1874; $\log_2 R = 1.2$ to 1.5), and ATP-dependent protease La (DVU3303) were found among other highly up-regulated genes (Table 2).

The *D. vulgaris* gene coding for the putative Na^+/H^+ antiporter NhaC-2 (DVU3108) is of particular interest. The role of Na^+/H^+ antiporters in regulation of intracellular pH homeostasis and survival at high pH has been demonstrated for a number of bacteria (5, 16, 28, 29). Although several genes in *D. vulgaris* have been identified as putative Na^+/H^+ antiporters,

TABLE 2. Selected up-regulated genes during pH 10 stress

Gene	Description	Expression change (log ₂ ratio) at ^a :			
		30 min	60 min	120 min	240 min
DVU0258	Sensory transduction histidine kinase related	0.96	0.95	1.24	2.28
DVU0260	Response regulator	0.96	1.30	1.23	1.25
DVU0261	Universal stress protein family	0.59	0.73	0.48	0.31
DVU0262	Hypothetical protein	0.53	1.25	0.95	1.29
DVU0263	Tetraheme cytochrome <i>c</i> ₃	0.22	0.41	0.20	0.69
DVU0264	Ferredoxin, 2[4Fe-4S]	0.79	0.77	0.94	0.88
DVU0265	25.3-kDa protein in <i>hmc</i> operon	1.61	2.08	1.54	1.56
DVU0331	Putative histidine protein kinase	1.36	1.18	1.83	2.23
DVU0667	HD domain protein	2.28	1.52	1.74	1.69
DVU0692	Molybdopterin oxidoreductase	1.09	1.21	1.52	2.01
DVU0693	Respiratory nitrate reductase, beta subunit (<i>narH</i>)	1.09	1.52	1.48	2.01
DVU0694	Molybdopterin oxidoreductase, molybdopterin binding subunit	2.17	1.91	1.01	1.97
DVU0774	ATP synthase F1, epsilon subunit (<i>atpC</i>)	1.77	2.57	2.20	3.21
DVU0775	ATP synthase F1, beta subunit (<i>atpD</i>)	1.11	1.55	1.63	2.27
DVU0776	ATP synthase F1, gamma subunit (<i>atpG</i>)	0.87	1.60	1.13	2.26
DVU0777	ATP synthase F1, alpha subunit	0.38	0.67	0.97	1.49
DVU0778	ATP synthase F1, delta subunit	0.77	1.03	1.31	1.86
DVU0779	ATP synthase F0, B subunit	0.20	0.27	0.49	1.48
DVU0780	ATP synthase F0, B subunit	0.41	0.23	0.28	1.02
DVU1030	Universal stress protein family	1.19	0.97	1.14	1.64
DVU1035	Glucokinase (<i>glk</i>)	1.09	2.03	2.11	1.82
DVU1198	Riboflavin synthase, beta subunit (<i>ribH</i>)	1.94	2.05	1.59	2.03
DVU1446	Heptosyltransferase family	2.98	2.30	2.56	1.63
DVU2367	Acyl-(acyl carrier protein)-UDP- <i>N</i> -acetylglucosamine <i>O</i> -acyl transferase	0.74	0.73	0.97	0.26
DVU2368	(3- <i>R</i>)-Hydroxymyristoyl-(acyl carrier protein) dehydratase (<i>fabZ</i>)	1.64	1.90	1.87	2.42
DVU2369	UDP-3- <i>O</i> -(<i>R</i> -3-hydroxymyristoyl)-glucosamine <i>N</i> -acyltransferase	0.78	0.79	1.07	1.07
DVU2370	Outer membrane protein OmpH, putative	1.58	1.80	1.62	1.64
DVU2526	Periplasmic (NiFe) hydrogenase, large subunit	1.68	1.24	1.42	1.98
DVU2571	Ferrous iron transport protein B	1.15	1.42	1.53	1.20
DVU2572	Ferrous iron transport protein A, putative	1.25	1.36	1.80	1.76
DVU2816	Efflux system protein	1.04	1.80	1.87	1.95
DVU3025	Pyruvate-ferredoxin oxidoreductase	0.64	0.54	0.70	1.19
DVU3026	L-Lactate permease family protein	0.01	0.10	0.18	0.59
DVU3027	Glycolate/lactate oxidase, subunit GlcD	0.88	0.73	0.93	1.25
DVU3028	Glycolate oxidase, iron-sulfur subunit	1.57	2.07	1.92	2.30
DVU3029	Phosphate acetyltransferase	0.13	0.13	0.13	0.46
DVU3030	Acetate kinase	0.45	0.40	0.51	0.62
DVU3035	Methyl-accepting chemotaxis protein, putative	1.41	1.74	1.31	1.56
DVU3081	Integral membrane protein, putative	0.27	2.38	1.35	2.64
DVU3108	Na ⁺ /H ⁺ antiporter NhaC (<i>nhaC</i>)	1.36	1.60	1.41	2.10
DVU3110	L-Aspartate oxidase, putative	0.91	1.19	1.35	2.45
DVU3298	Hypothetical protein	1.38	1.86	1.09	2.15
DVU3299	Membrane protein	0.67	0.74	1.73	1.76
DVU3300	Hypothetical protein	0.68	2.27	0.84	3.57
DVU3301	Hypothetical protein	0.88	3.36	1.53	3.70
DVU3303	ATP-dependent protease La, putative	0.87	0.21	1.03	1.97
DVU3304	Sensor histidine kinase	0.62	0.41	0.66	1.11

^a Log₂ gene expression ratios at different times after the pH was changed compared with the control at the same times.

only the expression of *nhaC-2* changed substantially in response to high pH. Our analysis of an *nhaC-2* deletion mutant demonstrated that there was increased sensitivity to pH 8.9 relative to the wild type, which was accentuated at higher concentrations of sodium chloride (Fig. 2 and 3). Increased cell lysis in stationary phase was observed for this mutant but not for the wild type (Fig. 3B and D). This phenotype is similar to that of an *E. coli nhaA*-deficient mutant (29).

The DVU3110 gene located in close proximity to *nhaC-2* and transcribed in the same direction was also highly up-regulated at high pH. This gene was predicted to encode a putative flavin adenine dinucleotide-binding oxidoreductase/L-aspartate oxidase with a bona fide homolog present in only one other *Desulfovibrio* strain (*D. vulgaris* DP4; >99% identity;

GenBank accession number CP000527.1). The next most closely related amino acid sequence (42.8% identity) was found in the genome of *Magnetospirillum magneticum* AMB-1 (GenBank accession number AP007255.1), annotated as a member of the succinate dehydrogenase/fumarate reductase protein family (COG1053). The absence of genes typically associated with characterized succinate dehydrogenases and fumarate reductases, such as genes for *b*-type cytochromes and small membrane anchor proteins in the vicinity of DVU3110, does not allow us to suggest that DVU3110 codes for a protein with similar activities. In addition, neither fumarate nor succinate was detected in cultural medium of *D. vulgaris* exposed to pH 10 (data not shown). Nonetheless, the presence of a gene encoding 4Fe-4S ferredoxin (DVU3109) immediately down-




FIG. 2. Growth characteristics of selected *D. vulgaris* mutants at elevated pH with 50 mM glycine buffer. (A) pH 7.0. (B) pH 7.9. (C) pH 8.9. The data are the means of triplicate cultures, and the observed variability was within 10%. WT, wild type.

stream of DVU3110 suggests that the latter might encode for an oxidoreductase. An identical gene context surrounds the DVU3108 homolog in *D. vulgaris* DP4. Since an ortholog of DVU3110 has not been identified in the sequenced genome of another *Desulfovibrio* species (*D. desulfuricans* G20; GenBank accession number CP000112.1), a response to elevated pH via the combined activities of the oxidoreductase/L-aspartate oxidase and NhaC may be a species-specific strategy. Of additional note, genes identified as L-aspartate oxidase genes in the genomes of many *Gammaproteobacteria* (for instance, *E. coli*, *Nitrosococcus oceanii*, and *Methylococcus capsulatus*) are immediately upstream of a gene for a putative sigma E, suggesting that there is more general involvement of this type of reductase/L-aspartate oxidase in the stress response.

In addition to increased transcription of the gene encoding L-aspartate oxidase, a number of genes involved in amino acid synthesis and metabolism were also consistently up-regulated.

Genes encoding tryptophan synthase subunits α and β (DVU0471 and DVU0470) and other members of the tryptophan operon (DVU0460 to DVU0469) were slightly or moderately up-regulated (Table 3). In addition, transcription of a gene annotated as a tryptophanase gene (DVU2204) was elevated after 240 min of pH stress ($\log_2 R = 1.3$). The orthologous gene in *E. coli* (*tnaA*) was previously shown to be up-regulated at pH 9 (25, 34), and growth of a *D. vulgaris* mutant with this gene deleted was diminished at pH 8 (data not shown). The concerted increase in transcription of genes for biosynthesis and transport of amino acids other than tryptophan (genes for cysteine synthase A, dihydrodipicolinate reductase involved in lysine biosynthesis, isopropylmalate dehydratase involved in leucine biosynthesis, and homoserine dehydrogenase involved in aspartate biosynthesis) suggests that *D. vulgaris* employs multiple components of amino acid metabolism for survival at high pH (Table 3).




FIG. 3. Growth characteristics of selected *D. vulgaris* mutants at elevated pH and sodium chloride concentrations with 50 mM Tris buffer. (A) pH 7.0 and 40 mM NaCl. (B) pH 7.9 and 40 mM NaCl. (C) pH 7.0 and 80 mM NaCl. (D) pH 7.9 and 80 mM NaCl. The data are the means of triplicate cultures, and the observed variability was within 10%. wt, wild type.

TABLE 3. Gene expression profile of selected genes involved in amino acid metabolism

Gene	Description	Expression change (log ₂ ratio) at ^a :			
		30 min	60 min	120 min	240 min
DVU0085	Tryptophan synthase, beta subunit	1.36	1.75	1.90	2.45
DVU0086	Hypothetical protein	1.18	1.88	0.84	3.00
DVU0285	Imidazole glycerol phosphate synthase	0.88	1.03	1.10	0.15
DVU0286	Imidazole glycerol phosphate synthase, cyclase subunit	1.79	1.39	1.53	0.66
DVU0339	D-Isomer-specific 2-hydroxyacid dehydrogenase family	1.01	0.85	1.20	1.24
DVU0460	Predicted phospho-2-dehydro-3-deoxyheptonate aldolase	-0.08	0.04	0.15	0.57
DVU0461	Predicted 3-dehydroquininate synthase	0.36	0.11	0.54	0.42
DVU0462	Chorismate mutase/prephenate dehydratase	-0.04	0.26	0.73	0.47
DVU0463	3-Phosphoshikimate 1-carboxyvinyltransferase	0.52	0.34	0.93	0.46
DVU0464	Prephenate dehydrogenase	-0.04	0.12	0.56	1.02
DVU0465	Anthranilate synthase, component I	0.87	0.20	0.51	0.48
DVU0466	Anthranilate synthase, glutamine amidotransferase component	0.18	0.37	0.08	0.07
DVU0467	Anthranilate phosphoribosyltransferase	1.16	0.70	1.18	0.97
DVU0468	Indole-3-glycerol phosphate synthase	1.01	0.85	1.41	1.32
DVU0469	N-(5-Phosphoribosyl)anthranilate isomerase	0.72	0.93	1.27	0.99
DVU0470	Tryptophan synthase, beta subunit	0.61	0.66	0.71	0.74
DVU0471	Tryptophan synthase, alpha subunit	0.99	1.16	1.15	1.31
DVU0663	Cysteine synthase A	0.83	1.31	1.65	0.82
DVU0890	Homoserine dehydrogenase	1.33	1.55	1.69	1.19
DVU1466	Acetylglutamate kinase	1.43	1.01	1.36	1.04
DVU1585	Vitamin B ₁₂ -dependent methionine synthase family protein	1.53	1.66	1.27	1.00
DVU1609	Dihydrodipicolinate reductase	1.31	1.27	1.13	0.73
DVU1610	Glutamine-dependent NAD ⁺ synthetase	1.51	1.09	1.52	1.37
DVU2981	2-Isopropylmalate synthase	0.49	0.65	0.48	0.34
DVU2982	3-Isopropylmalate dehydratase, large subunit, putative	0.89	0.87	1.11	0.70
DVU2983	3-Isopropylmalate dehydratase, small subunit	0.60	0.76	1.03	0.30
DVU2984	Conserved hypothetical protein	0.96	0.97	1.30	1.00
DVU3048	Aspartate semialdehyde dehydrogenase	1.09	1.30	1.73	1.50
DVU3371	5-Methyl-homocysteine S-methyltransferase	0.54	1.55	2.18	1.87

^a Log₂ gene expression ratios at different times after the pH was changed compared with the control at the same times.

The integrity of the cell envelope is also challenged at elevated pH. Changes in the expression of several genes involved in cell wall and membrane biogenesis were differentially expressed following an increase in the pH of the growth medium. Increased transcription of a gene encoding the heptosyltransferase family protein (DVU1446), involved in the synthesis of the inner core region of lipopolysaccharide, was a notable example (Table 2). One of the most highly up-regulated genes associated with cell envelope structure was *fabZ* (DVU2368), encoding a putative beta-hydroxyacyl-(acyl carrier protein) dehydratase. This gene, involved in fatty acid biosynthesis, is part of a predicted six-gene operon containing, in addition to *fabZ*, genes encoding UDP-*N*-acetylglucosamine *O*-acyltransferase (DVU2367), UDP-3-*O*-(*R*-3-hydroxymyristoyl)-glucosamine *N*-acyltransferase (DVU2369), outer membrane protein OmpH (DVU2370), *N*-acetylmuramoyl-L-alanine amidase (DVU2371), and a hypothetical protein (DVU2372). The first two enzymes are involved in lipopolysaccharide biosynthesis, while the latter two are part of the peptidoglycan biosynthetic pathway. Transcription of the other three genes in this operon (DVU2367, DVU2369, and DVU2370) was also increased at pH 10 (the log₂ *R* values were 0.8 to 1.0, 0.78 to 1.07, and 1.6 to 1.8, respectively). Transcription of two genes (DVU2371 and DVU2372) located upstream and separated from the genes mentioned above by 151 bp was almost unchanged, suggesting that they do not belong to the same operon. The *fabZ* gene in *E. coli* is part of the sigma E regulon, which includes a sigma E consensus promoter, and is up-regulated following heat shock (30). Al-

though the expression of this gene in *E. coli* was reported not to be altered by either an increase or a decrease in pH, in *D. vulgaris* the transcription of all genes in the operon containing the *fabZ* ortholog increased following the shift to higher pH. Changes in the expression of these genes in *D. vulgaris* have not been detected in response to other environmental stressors, including salt, nitrate, potassium chloride, and heat shock (6, 13, 27; J. D. Wall, unpublished data).

Multiple genes associated with energy generation and electron transfer reactions showed increased expression at elevated pH. For instance, DVU0692, DVU0693, and DVU0694 were significantly up-regulated after 30 min of pH stress. These genes were previously annotated as genes encoding subunits of a molybdopterin oxidoreductase with unknown specificity. Analysis of amino acid sequences predicted that the proteins encoded by DVU0694 (the first gene in the operon) and DVU0692 contain 2 and 10 transmembrane helices, respectively. Because they both also contain putative signal peptide sequences, it is likely that together they comprise a membrane-bound protein complex. The transcription of several other genes predicted to be involved in energy generation and electron transfer reactions also increased. These genes included genes encoding the large subunit of periplasmic NiFe hydrogenase isozyme 2 (DVU2526) and the Fe-S subunit of glycolate/lactate oxidase (DVU3028). In addition, genes for formate dehydrogenases (DVU0587, DVU0588, DVU2481, and DVU2482), thiosulfate reductase (DVU0179), and CO-induced hydrogenase (DVU2286 to DVU2291) were moderately




FIG. 4. Most highly expressed operons during pH stress, including genes encoding a putative Lon protease (DVU3303), two regulators (DVU3304 and DVU3305), and set of membrane proteins of unknown function (DVU3299 to DVU3301). Open arrows indicate genes whose expression was up-regulated; gray arrows indicate genes whose expression was unchanged. Abbreviations for domains for the Lon protease: Lon, Lon protease domain; REC, receiver domain; AAA, ATPase domain; Lon C, Lon protease C-terminal proteolytic domain. prts, proteins.

up-regulated. Together with the fact that lactate consumption and acetate production were inhibited, these trends suggested that there was redirection of electron flow from sulfate reduction to unknown electron acceptors, possibly to maintain cytoplasmic redox status during the stress.

Many genes showing differential expression during alkaline pH stress code for proteins of unknown function for which BLAST analysis did not reveal any homologous proteins in publicly available databases. Thus, their possible role in surviving at high pH is obscure. Among the genes most highly up-regulated following 240 min at pH 10 were DVU3300 and DVU3301 ($\log_2 R = 3.70$ and $\log_2 R = 3.57$, respectively). These genes most likely comprise an operon containing two additional genes, DVU3298 and DVU3299, that were also up-regulated at alkaline pH (Fig. 4). Amino acid sequence analysis predicted that the proteins encoded by these genes possess at least one transmembrane segment and a signal peptide, suggesting that there is a membrane association. These genes were also previously shown to be up-regulated in *D. vulgaris* in response to acid, nitrate, nitrite, and sodium chloride stresses (13, 27; Wall, unpublished data).

Changes in expression of signal transduction and regulatory genes. The effect of high pH on genes categorized as signal transduction genes revealed that of the 273 genes in this category, 87 showed significant differential expression for at least one time interval, as judged by a *Z* score greater than 1.5. The *Z* score was greater than 2 for 44 of them. A peculiar feature of the *D. vulgaris* genome is the relatively large number of open reading frames annotated as genes encoding methyl-accepting chemotaxis proteins (MCPs). For example, *E. coli* has 5 MCP genes, compared with 27 MCP genes in *D. vulgaris*. In *E. coli* these proteins are involved in chemotactic responses, transducing signals to the flagellar motor. Three *D. vulgaris* MCP genes were up-regulated at pH 10. One of these genes (DVU3035) is located in close proximity to an operon comprised of genes presumably involved in assimilation of lactate and its conversion to acetate, a central metabolic pathway in *D. vulgaris*. This gene was previously shown to be up-regulated with salt stress (27). A second gene (DVU3082) was up-regulated twofold after 120 and 240 min of alkaline pH exposure. This gene was previously shown to be up-regulated with acid stress but to be unaffected by other stressors examined so far for *D. vulgaris*.

The third gene (DVU1884), up-regulated twofold after 240 min of alkaline pH, was previously shown to be also up-regulated with heat shock and nitrate stress (6, 13). Interestingly, the latter gene is immediately downstream of a locus consisting of 14 genes (including genes encoding ClpB, DnaJ, and peptidyl-prolyl *cis-trans* isomerase) associated with the *E. coli* general stress response. Although expression of flagellar genes in *D. vulgaris* was decreased and the motility of the cells was suppressed, the MCPs may be involved in regulation of cellular responses other than motility. For instance, it was recently demonstrated that a chemosensory-like pathway regulates developmental gene expression in *Myxococcus xanthus* (18).

The DVU0667 gene coding for an HD domain protein was among the most highly up-regulated genes in cells exposed to pH 10 for 30 min. Proteins with the HD motif belong to a superfamily of metal-dependent phosphohydrolases that include a variety of uncharacterized proteins associated with nucleotidyltransferases and helicases from *Bacteria*, *Archaea*, and *Eukaryotes*. Increases in expression of these genes suggest that *D. vulgaris* may respond to high pH by activating unidentified regulatory pathways that affect RNA modifications and stability.

There are only a few genes encoding alternative sigma factors in *D. vulgaris* (sigma N, sigma H, and a flagellar sigma factor). As reported by Chhabra et al. (6), neither a gene for sigma E nor a gene for RpoS has been found in the genome of this bacterium. This suggests that *D. vulgaris* may respond to stresses that cause misfolding or degradation of cell envelope proteins differently than better-studied bacteria. Indeed, in addition to the absence of sigma E, no *deg* and *rse* orthologs have been recognized in the genome. These genes have been shown to play an important role in the response of *E. coli* to periplasmic stress (30, 31). Thus, *D. vulgaris* apparently uses a system distinct from that of *E. coli* to control expression of genes encoding proteases and chaperone-like proteins. One of the most highly up-regulated genes (DVU3303) encodes a protease that is homologous to Lon but has a very peculiar domain structure. This protease contains a signal receiver domain common to CheY, OmpR, NtrC, and PhoB, as well as a phosphor acceptor site for histidine kinase homologs. Its carboxyl terminus contains ATPase and Lon protease (S16) proteolytic domains (Fig. 4). This domain structure suggests that this protease can be activated by phosphorylation and transfers the signal to an unknown protein which might be subsequently functionally altered by cleavage. Immediately downstream of DVU3303 are genes for a histidine kinase (DVU3304) and a putative response regulator (DVU3305), making this gene cluster a plausible candidate for a regulatory circuit involved in the response to pH stress and other stresses, since these regulatory genes were shown to be also up-regulated under pH 5.5, nitrate, chromium, or sodium chloride stress (13, 27, J. D. Wall, unpublished data).

Two additional genes annotated as genes encoding histidine kinases (DVU0331 and DVU2580) were also up-regulated in response to elevated pH. Mutants with mutations in these two genes, JW3011 and JW3024, demonstrated increased sensitivity to elevated pH (Fig. 2). Since the genes encoding these kinases are proximal to genes for cell wall biogenesis, some of which were down-regulated, they may also be part of a more general *D. vulgaris* response to cell envelope stress.




FIG. 5. Conceptual model of *D. vulgaris* response to high pH stress. The arrows next to protein, enzyme, and pathway names indicate changes in the corresponding gene expression.

Genes down-regulated during exposure to high pH. Genes for peptidyl-prolyl *cis-trans* isomerase (DVU2569; $\log_2 R = -1.4$ to -1.8), L-isoaspartate *O*-methyltransferase (DVU1849; $\log_2 R = -1.2$ to -1.4), and peptidyl-prolyl *cis-trans* isomerase B-2 (DVU1873; $\log_2 R = -0.8$ to -1.3) were found to be moderately down-regulated with alkaline stress. All three of these proteins are involved in cell wall biosynthesis, and observed changes in their expression may reflect a pause in cell growth and/or adaptive changes in cell wall composition to the stress. In addition, some energy production and central metabolism genes were consistently down-regulated, including genes encoding pyruvate carboxylase (DVU1834; $\log_2 R = -1.4$ to -1.8), B₁₂ binding domain protein/radical *S*-adenosylmethionine domain protein (DVU3016; $\log_2 R = -1.5$ to -2.1), desulfoferredoxin (DVU3183; $\log_2 R = -1.0$ to -1.2), ferredoxin II (DVU0305; $\log_2 R = -1.5$ to -1.9), and an L-lactate permease family protein (DVU2451; $\log_2 R = -1.1$ to -1.8). The *hupD* gene (DVU1923; $\log_2 R = -0.6$ to -1.6) for hydrogenase expression/formation was down-regulated after 240 min of pH 10 stress. Interestingly, 28 of the 57 genes most down-regulated during high-pH stress ($\log_2 R$ range, -1.5 to -3.6 ; average $\log_2 R$, ~ -2) were annotated as phage or transposon related, suggesting that control of their expression might be an important aspect of the *D. vulgaris* stress response or part of the phage survival strategy (see Table S3 in the supplemental material).

Conclusions. The model shown in Fig. 5 draws upon the data presented in this study and provides an overview of the response of *D. vulgaris* to alkaline stress. The observed increase in expression of genes for ATP synthase and the Na⁺/H⁺ antiporter NhaC-2 attenuates alkalization of the cytoplasm via increased proton import and is similar to the responses of *E. coli* and other bacteria (20, 25, 29). Similarly, a decrease in expression of flagellar genes and presumptive acidification of the cytoplasm via increased expression of genes involved in amino acid metabolism are also shared responses. Increased

expression of several genes for proteins potentially involved in energy generation or electron transfer reactions was also observed. These genes included genes for a formate hydrogenase, a molybdopterin oxidoreductase, cytoplasmic Coo hydrogenase, and a periplasmic NiFe hydrogenase. The latter changes may be part of a more general strategy to retain a cellular redox state necessary to sustain cellular functions when lactate consumption and sulfate respiration are repressed.

Although growth was arrested at pH 10, the cells remained metabolically active, as assessed by rescue in culture and a complex transcriptional response during the period of exposure. As part of this response there appeared to be a modification of the cell envelope, indicated by modulation of the expression of genes involved in cell wall biosynthesis. Some genes in this category showed decreased expression, whereas others were up-regulated. We anticipate that ongoing studies of the cellular proteome and membrane composition will better elucidate the character and possible function of any changes in membrane structure.

Finally, our observations suggest that although the response of *D. vulgaris* to increased pH is similar to the response of *E. coli*, this response is controlled by unique regulatory circuits. The alternative sigma factors (sigma S and sigma E) contributing to this stress response in *E. coli* are apparently absent in *D. vulgaris*. Our transcriptional analysis identified several regulators that are likely part of a *D. vulgaris*-specific stress response system. These regulators now provide specific targets for continued biochemical and genetic characterization of the stress response system of this organism and its relationship to regulatory pathways in other bacteria, particularly within the family *Desulfovibrionaceae*.

ACKNOWLEDGMENTS

We thank N. Pinel, S. Flagan, C. Walker, and K. Hillesland for careful reading of the manuscript and for valuable discussions and M. Sadilek for his invaluable assistance in performing the gas chromatog-

raphy-mass spectrometry and high-performance liquid chromatography analyses. We also thank J. Jacobsen for her assistance with gene expression data analysis and handling and J. A. Ringbauer, Jr., for generating mutants. The generosity of William W. Metcalf for providing the pRL27 plasmid for mutagenesis with the Tn5-RL27 transposon is appreciated.

This work was supported by the Department of Energy's Office of Biological and Environmental Sciences under the GTL-Genomics Program via the Virtual Institute for Microbial Stress and Survival (<http://VIMSS.lbl.gov>).

REFERENCES

- Abildgaard, L., M. B. Nielsen., K. U. Kjeldesen, and K. Ingvorsen. 2006. *Desulfovibrio alkalitolerans* sp. nov., a novel alkalitolerant, sulphate-reducing bacterium isolated from district heating water. *Int. J. Syst. Evol. Microbiol.* **56**:1019–1024.
- Aeckersberg, F., F. Bak, and F. Widdel. 1991. Anaerobic oxidation of saturated hydrocarbons to CO₂ by a new type of sulfate-reducing bacterium. *Arch. Microbiol.* **156**:5–14.
- Beller, H. R., M. Reinhard, and D. Grbić-Galić. 1992. Metabolic by-products of anaerobic toluene degradation by sulfate-reducing enrichment cultures. *Appl. Environ. Microbiol.* **58**:3192–3195.
- Bender, K. S., H.-C. B. Yen, C. L. Hemme, Z. Yang, Z. He, J. Zhou, T. C. Hazen, K. H. Huang, E. J. Alm, A. P. Arkin, and J. D. Wall. 2007. Analysis of a ferric uptake regulator (Fur) mutant of *Desulfovibrio vulgaris* Hildenborough. *Appl. Environ. Microbiol.* **73**:5389–5400.
- Cheng, J., A. A. Guffanti, and T. A. Krulwich. 1994. The chromosomal tetracycline resistance locus of *B. subtilis* encodes a Na⁺/H⁺ antiporter that is physiologically important at elevated pH. *J. Biol. Chem.* **269**:27365–27371.
- Chhabra, S. R., Q. He, K. H. Huang, S. P. Gaucher, E. J. Alm, Z. He, M. Z. Hadi, T. C. Hazen, J. D. Wall, J. Zhou, A. P. Arkin, and A. K. Singh. 2006. Global analysis of heat shock response in *Desulfovibrio vulgaris* Hildenborough. *J. Bacteriol.* **188**:1817–1828.
- Chun, K. T., H. J. Edenberg, M. R. Kelley, and M. G. Goebel. 1997. Rapid amplification of uncharacterized transposon-tagged DNA sequences from genomic DNA. *Yeast* **13**:233–240.
- DeWeerd, K. A., F. Concannon, and J. M. Sufita. 1991. Relationship between hydrogen consumption, dehalogenation, and the reduction of sulfur oxyanions by *Desulfomonile tiedjei*. *Appl. Environ. Microbiol.* **57**:1929–1934.
- Dilworth, M. J., and A. R. Glenn. 1999. Problems of adverse pH and bacterial strategies to combat it. *Novartis Found. Symp.* **221**:4–14.
- Dyrløv, J. B., H. Nielsen, G. von Heijne, and S. Brunak. 2004. Improved prediction of signal peptides: SignalP 3.0. *J. Mol. Biol.* **340**:783–795.
- Fry, N. K., J. K. Fredrickson, S. Fishbain, M. Wagner, and D. A. Stahl. 1997. Population structure of microbial communities associated with two deep, anaerobic, alkaline aquifers. *Appl. Environ. Microbiol.* **63**:1498–1504.
- Fu, R., and G. Voordouw. 1997. Targeted gene-replacement mutagenesis of *dcrA*, encoding an oxygen sensor of the sulfate-reducing bacterium *Desulfovibrio vulgaris* Hildenborough. *Microbiology* **143**:1815–1826.
- He, Q., K. H. Huang, Z. He, E. J. Alm, M. W. Fields, T. C. Hazen, A. P. Arkin, J. D. Wall, and J. Zhou. 2006. Energetic consequences of nitrite stress in *Desulfovibrio vulgaris* Hildenborough, inferred from global transcriptional analysis. *Appl. Environ. Microbiol.* **72**:4370–4381.
- Heidelberg, J. F., R. Seshadri, S. A. Haveman, C. L. Hemme, I. T. Paulsen, J. F. Kolonay, J. A. Eisen, N. Ward, B. Methe, L. M. Brinkac, S. C. Daugherty, R. T. DeJoy, R. J. Dodson, A. S. Durkin, R. Madupu, W. C. Nelson, S. A. Sullivan, D. Fouts, D. H. Haft, J. Selengut, J. D. Peterson, T. M. Davidsen, N. Zafar, L. Zhou, D. Radune, G. Dimitrov, M. Hance, K. Tran, H. Khouri, J. Gill, T. R. Utterback, T. V. Feldblyum, J. D. Wall, G. Voordouw, and C. M. Fraser. 2004. The genome sequence of the anaerobic, sulfate-reducing bacterium *Desulfovibrio vulgaris* Hildenborough. *Nat. Biotechnol.* **22**:554–559.
- Hofmann, K., and W. Stoffel. 1993. TMbase—a database of membrane spanning proteins segments. *Biol. Chem.* **374**:166–168.
- Ito, M., A. A. Guffanti, B. Oudega, and T. A. Krulwich. 1999. *mmp*, a multi-gene, multifunctional locus in *Bacillus subtilis* with roles in resistance to cholate and to Na⁺ and pH homeostasis. *J. Bacteriol.* **181**:2394–2402.
- Juncker, A. S., H. Willenbrock, G. von Heijne, H. Nielsen, S. Brunak, and A. Krogh. 2003. Prediction of lipoprotein signal peptides in Gram-negative bacteria. *Protein Sci.* **12**:1652–1662.
- Kirby, J. R., and D. R. Zusman. 2003. Chemosensory regulation of developmental gene expression in *Myxococcus xanthus*. *Proc. Natl. Acad. Sci. USA* **100**:2008–2013.
- Larsen, R. A., M. M. Wilson, A. M. Guss, and W. W. Metcalf. 2002. Genetic analysis of pigment biosynthesis in *Xanthobacter autotrophicus* Py2 using a new, highly efficient transposon mutagenesis system that is functional in a wide variety of bacteria. *Arch. Microbiol.* **178**:193–201.
- Leapheart, A. B., D. K. Thompson, K. Huang, E. Alm, X. Wan, A. P. Arkin, S. D. Brown, L. Wu, T. Yan, X. Liu, G. S. Wickman, and J. Zhou. 2006. Transcriptome profiling of *Shewanella oneidensis* gene expression following exposure to acidic and alkaline pH. *J. Bacteriol.* **188**:1633–1642.
- Lloyd, J. R., J. Ridley, T. Khiznia, N. N. Lyalikova, and L. E. Macaskie. 1999. Reduction of technetium by *Desulfovibrio desulfuricans*: biocatalyst characterization and use in a flowthrough bioreactor. *Appl. Environ. Microbiol.* **65**:2691–2696.
- Londry, K. L., J. M. Sufita, and R. S. Tanner. 1999. Cresol metabolism by the sulfate-reducing bacterium *Desulfotomaculum* sp. strain Groll. *Can. J. Microbiol.* **45**:458–463.
- Lovley, D. 1995. Bioremediation of organic and metal contaminants with dissimilatory metal reduction. *J. Ind. Microbiol.* **14**:85–93.
- Lovley, D. R., and E. J. Phillips. 1992. Reduction of uranium by *Desulfovibrio desulfuricans*. *Appl. Environ. Microbiol.* **58**:850–856.
- Maurer, L., E. Yohannes, S. S. Bondurant, M. Radmacher, and J. L. Slonczewski. 2005. pH regulates genes for flagellar motility, catabolism, and oxidative stress in *Escherichia coli* K-12. *J. Bacteriol.* **187**:304–319.
- Motamedi, M., and K. Pedersen. 1998. *Desulfovibrio aespoensis* sp. nov., a mesophilic sulfate-reducing bacterium from deep groundwater at Aspo hard rock laboratory, Sweden. *Int. J. Syst. Bacteriol.* **48**:311–312.
- Mukhopadhyay, A., Z. He, E. J. Alm, A. P. Arkin, E. E. Baidoo, S. C. Borglin, W. Chen, T. C. Hazen, Q. He, H. Y. Holman, K. Huang, R. Huang, D. C. Joyner, N. Katz, M. Keller, P. Oeller, A. Redding, J. Sun, J. Wall, J. Wei, Z. Yang, H. C. Yen, J. Zhou, and J. D. Keasling. 2006. Salt stress in *Desulfovibrio vulgaris* Hildenborough: an integrated genomics approach. *J. Bacteriol.* **188**:4068–4078.
- Padan, E., N. Maisler, D. Taglicht, R. Karpel, and S. Schuldiner. 1989. Deletion of *ant* in *E. coli* reveals its function in adaptation to high salinity and an alternative Na/H antiporter system(s). *J. Biol. Chem.* **264**:20297–20302.
- Padan, E., E. Bibi, M. Ito., and T. Krulwich. 2005. Alkaline pH homeostasis in bacteria: new insights. *Biochim. Biophys. Acta* **1717**:67–88.
- Rhodijs, V. A., W. C. Suh, G. Nanaka, and C. A. Gross. 2006. Conserved and variable functions of the sigma E stress response in related genomes. *PLoS Biol.* **4**:43–59.
- Ruiz, N., and T. Silhavy. 2005. Sensing external stress: watchdogs of the *E. coli* cell envelope. *Curr. Opin. Microbiol.* **8**:122–126.
- Saeed, A. I., V. Sharov, J. White, J. Li, W. Liang, N. Bhagabati, J. Braisted, M. Klapa, T. Currier, M. Thiagarajan, A. Sturn, M. Snuffin, A. Rezantsev, D. Popov, A. Ryltsov, E. Kostukovich, I. Borisovsky, Z. Liu, A. Vinsavich, V. Trush, and J. Quackenbush. 2003. TM4: a free, open-source system for microarray data management and analysis. *BioTechniques* **34**:374–378.
- Skulachev, V. P. 1999. Bacterial energetics at high pH: what happens to the H⁺ cycle when the extracellular H⁺ concentration decreases? *Novartis Found. Symp.* **221**:200–213.
- Stancik, L. M., D. M. Stancik, B. Schmidt, D. M. Barnhart, Y. N. Yoncheva, and J. L. Slonczewski. 2002. pH-dependent expression of periplasmic proteins and amino acid catabolism in *Escherichia coli*. *J. Bacteriol.* **184**:4246–4258.
- Vandijken, V., C. Knoblauch, and B. B. Jørgensen. 2006. *Desulfovibrio frigidus* sp. nov. and *Desulfovibrio ferrireducens* sp. nov., psychrotolerant bacteria isolated from Arctic fjord sediments (Svalbard) with the ability to reduce Fe(III). *Int. J. Syst. Evol. Microbiol.* **56**:681–685.
- Zhang, W., D. E. Culley, J. C. Scholten, M. Hogan, L. Vitoritti, and F. J. Brockman. 2006. Global transcriptomic analysis of *Desulfovibrio vulgaris* on different electron donors. *Antonie Leeuwenhoek* **89**:221–237.
- Zhang, W., D. E. Culley, M. Hogan, L. Vitoritti, and F. J. Brockman. 2006. Oxidative stress and heat-shock responses in *Desulfovibrio vulgaris* by genome-wide transcriptomic analysis. *Antonie Leeuwenhoek* **90**:41–55.