

1 ***In situ* Bioreduction of Uranium (VI) to Submicromolar**
2 **Levels and Reoxidation by Dissolved Oxygen**

3 **Supporting Information**

4 WEI-MIN WU¹, JACK CARLEY², JIAN LUO¹, MATTHEW A GINDER-VOGEL³, ERICK
5 CARDENAS⁴, MARY BETH LEIGH⁴, CHAICHI HWANG⁵, SHELLY D. KELLY⁶, CHAUNGMIN
6 RUAN², LIYOU WU⁷, JOY VAN NOSTRAND⁷, TERRY GENTRY², KENNETH LOWE², TONIA
7 MEHLHORN², SUE CARROLL², WENSUI LOU², MATTHEW W. FIELDS⁵, BAOHUA GU²,
8 DAVID WATSON², KENNETH M. KEMNER⁶, TERENCE MARSH⁴, JAMES TIEDJE⁴, JIZHONG
9 ZHOU⁷, SCOTT FENDORF³, PETER K. KITANIDIS¹, PHILIP M. JARDINE², AND CRAIG S.
10 CRIDDLE^{1*}

11 ¹ Department of Civil and Environmental Engineering , Stanford University, Stanford, CA 94305.

12 ² Environmental Sciences Division, Oak Ridge National Laboratory, P.O. Box 2008, Oak Ridge, TN
13 37831.

14 ³ Department of Geological and Environmental Sciences, Stanford University, Stanford, CA 94305.

15 ⁴ Center for Microbial Ecology, Michigan State University, East Lansing, MI 48824-1325.

16 ⁵ Department of Microbiology, Miami University, Oxford, OH 45056.

17 ⁶ Biosciences Division, Argonne National Laboratory, Argonne, IL, 60439.

18 ⁷ University of Oklahoma, Norman, OK 73019

19

20

21

22 **Bacterial Community Analysis**

23 *A) Groundwater samples.*

24 Total bacterial genomic DNA was extracted from the filters as described previously (Hwang et al.,
25 2006). The SSU rRNA genes were amplified in a PTC-200 Peltier Thermal Cycler (MJ Research) with
26 the primer pair FD1 F (5' AGA GTT TGA TCC TGG CTC AG 3') and 1540R (5' AAG GAG GTG
27 ATC CAG CC 3'). The PCR reaction (25 ul) contained 12.5 µL BioMix (Bioline, Randolph, MA), 1µL
28 each primer, 9.5 µL sterilized Milli-Q water, 1µL purified DNA (5-10 ng). To minimize PCR-induced
29 artifacts, the optimal number of cycles was determined and five PCR reactions were combined prior to
30 cloning as described previously (Qui et al. 2001). The PCR parameters were as follows: 80°C for 1.5
31 min; 94°C for 2 min; 94°C for 30s; 58°C for 1 min; 72°C for 1 min; 20 cycles; 72°C for 7 min. An
32 aliquot of 5 µL of PCR product was run in a 0.8% agarose TAE gel stained with ethidium bromide to
33 evaluate the quality of the amplified fragment.

34 PCR products of the SSU rRNA genes were purified using a Promega Kit (Promega, Madison,
35 WI). The purified fragments were cloned using the vector PCR 2.1-TOPO and *Escherichia coli*
36 DH5_ϕplainTM-T1^R competent cells according to the manufacturer's instructions (Invitrogen, Carlsbad,
37 CA). From each of the libraries approximately 100 white colonies were randomly selected, and the
38 cloned inserts were amplified with vector-specific primers M13 forward and M13 reverse (30 cycles of
39 the PCR parameters described above, but with an annealing temperature of 60°C and a final extension
40 time of 10 minutes). The resulting amplification products were analyzed as described above. PCR
41 products (20µl) amplified with vector-specific primers were purified with a Montage PCR_{u96} plate
42 according to the manufacturer's instructions (Millipore, Bedford, MA). DNA sequences were
43 determined with a DYEnamic ET Terminator Cycle Sequencing Kit (Amersham Biosciences,
44 Piscataway, NJ) according to manufacturer's instructions at 1:4 dilutions with an internal primer (529r).
45 Sequence reaction products were analyzed on the ABI model 3100 and 3730 DNA sequencer (Applied
46 Biosystems). DNA sequences were assembled and edited using the SequencherTM program (v. 4.0,
47 Gene Codes Corporation, Ann Arbor, MI). The SSU rDNA sequence identification was performed with
48 the BLASTn server (<http://www.ncbi.nlm.nih.gov/BLAST>) and the RDPII
49 (<http://rdp.cme.msu.edu/index.jsp>). Unique SSU rDNA clones from each library were identified and
50 designated as OTUs (operational taxonomic unit) at a 97% sequence identity level. The sequences were
51 aligned with Clustal W (Thompson et al. 1994). Phylogenetic trees were constructed with distance
52 matrices and the neighbor-joining method, and bootstrap analysis was performed with programs within

53 MEGA version 2.1 (Kumar et al. 2001).

54 ***B) Sediment Samples***

55 Sediments were collected from the wells in the treatment zone by a well surging operation, as described
56 previously (Wu et al., 2006b). Total bacterial DNA was extracted from 0.5 g of sediments with the Fast
57 DNA spin kit for soil (BIO 101, Carlsbad, CA) following the manufacturer's instructions.

58 16S rRNA genes were amplified using the universal primers 27F (5'-
59 AGAGTTTGATCMTGGCTCAF-3') and 1392R (5'-ACGGGCGGTGTGTRC-3'). Thermocycler
60 conditions were as follows: 95°C for 5 min; 95 °C for 1 min, 59 °C for 1 min, 72°C for 1:40 min, for 28
61 cycles; 72 °C for 10 min. PCR products were analyzed in a 1.5% agarose TAE gel and the size of the
62 product was confirmed. To control for PCR bias, four replicate PCR reactions were prepared for each
63 sample and then compiled. The combined PCR products (200 µl) were purified using a Qiaquick PCR
64 purification kit (Qiagen Inc., Valencia, CA, USA). The concentrated products were then run in a 1%
65 agarose gel, excised, and extracted with a Qiaquick gel extraction kit (Qiagen Inc., Valencia, CA).
66 Adenine overhangs were added to the PCR products by incubating a mixture of 18.8mM Tris HCl pH
67 8.4, 47mM KCl, 0.93mM dATP and 0.5 U of taq polymerase for 10 min at 72 °C. Prepared PCR
68 products were then cloned using the Topo TA Clonig kit for sequencing following manufacturer's
69 instructions (Invitrogen Life Technologies, Carlsbad, CA, USA). Single read sequencing was conducted
70 by Macrogen Inc (Seoul, Korea) using the internal primer 27F.

71 The raw sequence data (Ab1 files) were submitted to the Pipeline application of the Ribosomal
72 Database project (RDP at www.rdp.cme.msu.edu) (Cole et al., 2005, 2007). Bases were called using
73 PHRED (Ewing et al., 1998) and vector sequence removed with LUCY (Chou et al., 2001). The
74 resulting sequences were aligned to the RDP model and rRNA distances matrices were generated and
75 corrected using the Jukes – Cantor correction. Sequences were also downloaded with a Q20 quality
76 mask to be later classified using the RDP classifier. The sequences were classified with the classifier
77 application of the RDP using an 80% confidence value.

78 For methods section:

79 **Functional Gene Microarray**

80 A 50mer functional gene microarray containing 25,098 genes involved in biogeochemical cycling of C,
81 N, S, organic contaminant degradation, and metal homeostasis (Rhee et al., 2004; Li et al., 2005;
82 Liebich et al. 2006) was used to examine microbial communities from ground water. Samples were
83 obtained as described above. DNA was extracted from filters by the method of Zhou et al. (1996) and

84 50 ng aliquots of DNA were amplified, labeled and hybridized as described previously (Wu et al.,
85 2006).

86 **Most Probable Number (MPN) method**

87 Denitrifying bacteria, FeRB and SRB were enumerated using the Most Probable Number (MPN)
88 technique with five tubes for each dilution. Anaerobic pressure tubes (27 mL) containing 10 ml basal
89 medium were sealed with butyl rubber stoppers with aluminum caps. The basal medium contained the
90 following components (per liter): NH₄Cl, 0.5g; NaCl, 0.4g; NaHCO₃, 0.55g; and mineral solution, 100
91 ml. The mineral solution contained (per liter): MnCl₂, 0.4 g; MgSO₄, 1.5 g; CaCl₂, 0.5 g; and yeast
92 extract, 0.02g. The medium was prepared under a N₂-CO₂ (99:1, vol/vol) atmosphere and distributed to
93 each pressure tube (10 mL per tube). After autoclaving, a sterile trace element solution (0.4 ml) and a
94 sodium trimetaphosphate solution (50 mM, 0.025 ml) was added into each tube to obtain pH 7.0. The
95 trace element solution contained (per liter): HCl (12N), 6.4 mL; FeCl₂·4H₂O, 0.3g; ZnSO₄·H₂O, 0.1g;
96 MnSO₄, 0.085g; HB, 0.06g; CoCl₂·6H₂O, 0.02g; CuSO₄, 0.004g; NiSO₄·6H₂O, 0.028 g; and
97 NaMoO₃·2H₂O, 0.04g. The electron acceptor solution was supplemented to the tubes to obtain 5 mM
98 Fe-citrate for FeRB, 1.0 g/L of sodium thioglycollate and 5 g/l of FeSO₄ for SRB, and 1.0 g/L of NaNO₃
99 for denitrifiers, respectively. Ethanol solution (1 M) was added to each tube to give a final
100 concentration of 10 mM. Groundwater was pumped from the wells into anaerobic pressure tubes pre-
101 filled with nitrogen gas. The sample was then serially diluted in MPN tubes. The inoculated tubes were
102 incubated at ambient temperature for two months. Tubes were compared to controls for scoring as
103 positive or negative for production of gas in denitrifying tubes, color change in FeRB tubes, and
104 production of black Fe(II) precipitates in SRB tubes. FeRB and SRB tubes were also analyzed and
105 scored for methane production. Methane production in MPN tubes for FeRB and SRB was considered
106 indicative of the presence of methanogens.

107

108 **XANES Measurements at MR-CAT**

109 Some of the XANES analysis were carried out at Argonne National Laboratory (ANL) and
110 others at Stanford University. Samples measured at ANL are shown in Supplemental Figure 1. Sediment
111 samples from the wells were collected in anaerobic serum bottles with He or Ar headspace and stored at
112 4 °C in a refrigerator. They were centrifuged to separate the sediments from the supernatant within a
113 Coy(R) Anoxic Chamber filled with N₂ and H₂ mixture, mounted under anoxic conditions into
114 plexiglass sample holders with x-ray transparent windows of Kapton film covered with Kapton tape,
115 and stored at 4° C in canning jars filled with anoxic atmosphere for at most 3 days until measurements


116 were made at the MR-CAT (Segre et al., 2000) beamline at the Advance Photon Source, Argonne
117 National Laboratory, USA. The sealed samples were exposed to oxygen only briefly to place them in
118 the sample chamber at the beamline which was purged with N₂ gas. When exposed to atmospheric
119 conditions, sealed samples have been shown to preserve the anoxic integrity of the samples for more
120 than 24 hours. The average valence states of U within the moist sediments were measured under anoxic
121 conditions with no pretreatment of the sediments.

122 The MR-CAT insertion device beamline parameters were as follows. The insertion device was
123 operated on the third harmonic with a taper of approximately 2 keV to produce an incident x-ray
124 intensity as uniform as possible over the scanned x-ray region from 17,000 eV to 18,000 eV. A double
125 crystal Si(111) monochromator was used to select the x-ray energy. A Rh harmonic x-ray rejection
126 mirror was used to eliminate x-rays with higher harmonic energies. The x-ray profile at the sample was
127 approximately 0.7 mm square. The incident, transmitted, and fluorescence x-ray intensities were
128 measured with pure N₂, mixture of 90% N₂ and 10% Ar, and pure Ar filled ionization chambers,
129 respectively. The fluorescence ionization chamber was used with solar slits and an Al filter in the Stern-
130 Heald geometry (Stern et al, 1979; Kemner et al., 1994) performed with 50% attenuation in x-ray
131 intensity on reference samples indicated that the nonlinearity in the detectors was less than 0.5%. A
132 energy reference spectrum from a hydrogen uranyl phosphate standard was measured with each energy
133 scan. This reference was measured after the sample using a reference ionization chamber filled with a
134 mixture of 90% Ar and 10% N₂.

135 The monochromator was scanned continuously to collect quick spectra within ~ 2 minutes.
136 Long x-ray exposure times to the moist samples showed some changes in the XANES spectra.
137 Therefore two scans, resulting in less than 5% change in the spectra were collected from many different
138 regions of the uniform sample consisting of fine particulates surged from the wells. The spectra from
139 each sample were averaged to increase the signal to noise ratio within the final spectra. The averaged
140 spectra are shown in Supplemental Figure 1.

141 The XANES spectra were modeled with a linear combination of standard spectra with the
142 software Athena (Ravel et al., 2005) which uses IFEFFIT methods (Newville, et al., 2001). Two
143 different U(VI) and U(IV) spectra were used as possible standards as needed to reproduce the measured
144 spectra. The U(VI) standards are aqueous uranyl nitrate at low pH (denoted U(VI) Std-1) and a

145 hydrogen uranyl phosphate mineral (denoted U(VI) Std-2). The U(IV) standards are a natural uraninite
146 mineral (denoted U(IV) Std-1) and abiotically produced uraninite particles (denoted U(IV) Std-2)
147 (O'Loughlin, et al., 2000). These standards are shown in Supplemental Figure 1.
148



149
150 Supplemental Figure 1. XANES analysis of selected samples (black) and linear
151 combination fitting (red) with U(IV) and U(VI) standards (blue).
152

153 The linear combinations are shown with the measured spectra in Supplemental Figure 1. The
154 average U(IV) percentage is listed in Table 1 with the remaining percentage as U(VI). As shown in
155 Supplemental Figure 1, the XANES spectra depend largely on the U valence state but also to a lesser
156 extent on the U speciation. The reference spectra used in the linear combination fitting are from species
157 that are surely somewhat different from the actual species within the sediment samples, causing some
158 systematic error in the percentage of U(IV) to U(VI) within the sediment sample. Based on the

159 differences of the XANES spectra for different U(VI) and U(IV) species, we generously estimate the
160 accuracy of this method to be 10%.

161

162 **References**

163 Ewing, B.; Green, P. Base-calling of automated sequencer traces using phred. II. Error probabilities.
164 *Genome Research*, **1998**, 8:186-194.

165 Chou, H-H.; Holmes, M.H. DNA sequence quality trimming and vector removal. *Bioinformatics*. **2001**.
166 17: 1093-1104.

167

168 Cole, J.R.; Chai, B.; Farris, R.J.; Wang, Q.; Kulam, S.A.; McGarrell, D.M.; Garrity, G.M. and Tiedje,
169 J.M. The Ribosomal Database Project (RDP-II): sequences and tools for high-throughput rRNA
170 analysis. *Nucleic Acids Res.* **2005** January 1; 33(Database Issue): D294–D296.

171

172 Cole, J.R.; Chai, B.; Farris, R.J.; Wang, Q.; Kulam-Syed-Mohideen, A.S.; McGarrell, D.M.; Bandela,
173 A.M.; Cardenas, E.; Garrity, G.M; Tiedje, J.M. The Ribosomal Database Project (RDP-II): Introducing
174 myRDP space and quality controlled public data. *Nucleic Acids Res.* **2007**. (Submitted).

175

176 Hwang, C.; Wu, W.-M.; Gentry, T.; Carley, J.; Carroll, S.L.; Schadt, C.; Watson, D.; Jardine, P.M.;
177 Zhou, J.; Hickey, R.F.; Criddle, C.S.; Fields, M.W. Changes in bacterial community structure correlate
178 with initial operating conditions of a field-scale denitrifying fluidized bed reactor. *Appl. Microbiol.*
179 *Biotechnol.* **2006**, 71, 748-760.

180

181 Kemner, K. M.; Kropf, A. J.; Bunker, B. A. A low-temperature total electron yield detector for X-ray
182 absorption fine structure spectra. *Rev. Sci. Instrum.* **1994**, 65, 3667-3669.

183

184 Kumar, S.; Tamura, K.; Jakobsen, I.B.; Mei, M. MEGA 2: Molecular Evolutionary Genetics Analysis
185 Software. *Bioinformatics*. **2001**, 12, 1244-1245.

186 Li, X.; He, Z.; Zhou, J. Selection of optimal oligonucleotide probes for microarrays using multiple
187 criteria, global alignment and parameter estimation. *Nucleic Acids Res.* **2005**, 33, 6114-6123.

188

189 Liebich, J.; Schadt, C. W.; Chong, S. C.; He, Z.; Rhee, S. K.; Zhou, J. Improvement of oligonucleotide
190 probe design criteria for functional gene microarrays in environmental applications. *Appl. Environ.*
191 *Microbiol.* **2006**, 72, 1688-1691.

192

193 Newville, M. IFEFFIT: Interactive EXAFS analysis and FEFF fitting. *J. Synch. Rad.* **2001**, 8, 322-324.

194

195 O'Loughlin, E. J.; Kelly, S. D.; Cook, R. E.; Csencsits, R.; Kemner, K. M. Reduction of uranium(VI) by
196 mixed iron(II)/iron(III) hydroxide (green rust): Formation of UO₂ nanoparticles. *Environ. Sci. Technol.*
197 **2003**, 37, 721-727.

198

199 Qiu, X.; Wu, L.; Huang, H.; McDonel, P.E.; Palumbo, A.V.; Tiedje, J.M.; Zhou, J. Evaluation of PCR-
200 generated chimeras, mutations, and heteroduplexes with 16S rRNA gene-based cloning. *Appl. Environ.*
201 *Microbiol.* **2001**, 67:880-887.

202

203 Ravel, B.; Newville, M. ATHENA, ARTEMIS, HEPHAESTUS: Data analysis for X-ray absorption. *J.*
204 *Synch. Rad.* **2005**, 12, 537-541.

205

206 Rhee, S. K.; Liu, Z.; Wu, L.; Chong, S. C.; Wan, X.; Zhou, J. Detection of genes involved in
207 biodegradation and biotransformation in microbial communities by using 50-mer oligonucleotide

208 microarrays. *Appl. Environ. Microbiol.* **2004**, 70, 4303-4317.
209
210 Segre, C. U.; Leyarovska, N. E.; Chapman, L. D.; Lavender, W. M.; Plag, P. W.; King, A. S.; Kropf, A.
211 J.; Bunker, B. A.; Kemner, K. M.; Dutta, P.; Druan, R. S.; Kaduk, The MRCAT insertion device
212 beamline at the Advanced Photon Source. *J. Synchrotron Rad. Inst.* **2000**, CP521, 419-422.
213
214 Stern, E. A.; Heald, S. M. X-ray filter assembly for fluorescence measurements of x-ray absorption fine
215 structure. *Rev. Sci. Instrum.* **1979**, 50, 1579-1583.
216
217 Suzuki, Y.; Kelly, S. D.; Kemner, K. M.; Banfield, J. F. Enzymatic U(VI) reduction by
218 *Desulfosporosinus* species. *Radiochem. Acta.* **2004**, 92, 11-16.
219
220 Thompson, J.D.; Higgins, D.G.; Gibson, T.J. CLUSTAL W: improving the sensitivity of progressive
221 multiple sequence alignment through sequence weighting, positions-specific gap penalties and weigh
222 matrix choice. *Nucleic Acids. Res.* **1994**, 22, 4673-4680.
223 Wu, L.; Liu, X.; Schadt, C. W.; Zhou, J. Microarray-based analysis of subnanogram quantities of
224 microbial community DNAs by using whole-community genome amplification. *Appl. Environ.*
225 *Micrbiol.* **2006**, 72, 4931-4941.
226
227 Wu, Q.; Sanford, R. A.; Löffler, F. E. Uranium(VI) Reduction by *Anaeromyxobacter dehalogenans*
228 Strain 2CP-C. *Appl. Environ. Microbiol.* **2006**, 72, 3608-3614.
229
230 Zhou, J., Bruns, M. A., Tiegje, J. M. DNA recovery from soils of diverse composition *Appl. Environ.*
231 *Microbiol.* **1996**, 62, 316-322.