





PRIMARY RESEARCH ARTICLE

Long-term nitrogen addition modifies microbial composition and functions for slow carbon cycling and increased sequestration in tropical forest soil

Jing Tian^{1,2} | Jennifer A. J. Dungait³ | Xiankai Lu⁴ | Yunfeng Yang⁵  |
Iain P. Hartley³ | Wei Zhang⁴ | Jiangming Mo⁴  | Guirui Yu²  | Jizhong Zhou^{5,6,7} |
Yakov Kuzyakov^{8,9} 

¹College of Resources and Environmental Sciences, National Academy of Agriculture Green Development, Key Laboratory of Plant-Soil Interactions, Ministry of Education, China Agricultural University, Beijing, PR China

²Key Laboratory of Ecosystem Network Observation and Modeling, Institute of Geographic Sciences and Natural Resources Research, Chinese Academy of Sciences (CAS), Beijing, PR China

³Geography, College of Life and Environmental Sciences, University of Exeter, Exeter, UK

⁴Key Laboratory of Vegetation Restoration and Management of Degraded Ecosystems and Guangdong Provincial Key Laboratory of Applied Botany, South China Botanical Garden, Chinese Academy of Sciences, Guangzhou, China

⁵State Key Joint Laboratory of Environment Simulation and Pollution Control, School of Environment, Tsinghua University, Beijing, PR China

⁶Department of Microbiology and Plant Biology and School of Civil Engineering and Environmental Sciences, Institute for Environmental Genomics, University of Oklahoma, Norman, OK, USA

⁷Earth and Environmental Sciences, Lawrence Berkeley National Laboratory, Berkeley, CA, USA

⁸Department of Soil Science of Temperate Ecosystems, University of Göttingen, Göttingen, Germany

⁹Faculty of Life Science and Technology, Central South University of Forestry and Technology, Changsha, China

Correspondence

Jiangming Mo, Key Laboratory of Vegetation Restoration and Management of Degraded Ecosystems and Guangdong Provincial Key Laboratory of Applied Botany, South China Botanical Garden, Chinese Academy of Sciences, 510650 Guangzhou, China.
Email: mojm@scib.ac.cn

Guirui Yu, Key Laboratory of Ecosystem Network Observation and Modeling, Institute of Geographic Sciences and Natural Resources Research, Chinese Academy of Sciences (CAS), 100101 Beijing, PR China.
Email: yugr@igsrr.ac.cn

Funding information

National Natural Science Foundation of China, Grant/Award Number: 31770560, 41571130041 and 41731176; National Key R&D Program of China, Grant/Award Number: 2017YFA0604803; Youth Innovation Research, Grant/Award Number: LENOM2016Q0004

Abstract

Nitrogen (N) deposition is a component of global change that has considerable impact on belowground carbon (C) dynamics. Plant growth stimulation and alterations of fungal community composition and functions are the main mechanisms driving soil C gains following N deposition in N-limited temperate forests. In N-rich tropical forests, however, N deposition generally has minor effects on plant growth; consequently, C storage in soil may strongly depend on the microbial processes that drive litter and soil organic matter decomposition. Here, we investigated how microbial functions in old-growth tropical forest soil responded to 13 years of N addition at four rates: 0 (Control), 50 (Low-N), 100 (Medium-N), and 150 (High-N) kg N ha⁻¹ year⁻¹. Soil organic carbon (SOC) content increased under High-N, corresponding to a 33% decrease in CO₂ efflux, and reductions in relative abundances of bacteria as well as genes responsible for cellulose and chitin degradation. A 113% increase in N₂O emission was positively correlated with soil acidification and an increase in the relative abundances of denitrification genes (*narG* and *norB*). Soil acidification induced by N addition decreased available P concentrations, and was associated with reductions in the relative abundance of phytase. The decreased relative abundance of bacteria and key functional gene groups for C degradation were related to slower

SOC decomposition, indicating the key mechanisms driving SOC accumulation in the tropical forest soil subjected to High-N addition. However, changes in microbial functional groups associated with N and P cycling led to coincidentally large increases in N₂O emissions, and exacerbated soil P deficiency. These two factors partially offset the perceived beneficial effects of N addition on SOC storage in tropical forest soils. These findings suggest a potential to incorporate microbial community and functions into Earth system models considering their effects on greenhouse gas emission, biogeochemical processes, and biodiversity of tropical ecosystems.

KEYWORDS

biogeochemical cycling, C and N turnover, global climate change, microbial functional community, N deposition, tropical forest

1 | INTRODUCTION

Atmospheric deposition of reactive nitrogen (N) has increased significantly across every continent in the last few decades, with the most rapid increase observed in the northern tropical zone (Galloway et al., 2004; Tian et al., 2015). Investigations of the responses of temperate and boreal forest ecosystems to N deposition are relatively extensive compared to forests in the tropics, where most of them are generally of much shorter duration (Cusack et al., 2016; Pajares & Bohannan, 2016). N fertilization can increase carbon (C) sequestration in temperate and boreal forests by increasing net primary productivity (NPP) and slowing soil organic carbon (SOC) decomposition rates, but the net effect of N deposition on soil C storage in tropical forests remains unclear; this sink is extremely vulnerable to human perturbation, including the indirect effects of land use and climate change (Baccini et al., 2017; Cusack et al., 2016). Tropical forest soils are generally rich in N compared to the natural N-limitation in boreal and temperate ecosystems, and are the largest natural sources of N₂O emissions that are produced primarily by soil microbial nitrification/denitrification processes (Li, Niu, & Yu, 2016; Lu & Tian, 2013; Vitousek, Porder, Houlton, & Chadwick, 2010). In contrast, biologically available P is generally the limiting nutrient in old, highly weathered tropical forest soils (Camenzind, Hattenschwiler, Treseder, Lehman, & Rillig, 2018), and the effects of N addition on P cycling in these fragile environments are poorly understood. Consequently, the N deposition in tropical forest ecosystems may cause changes in soil microbial community composition and metabolic functions, reducing the potential for tropical forests to deliver globally important ecosystem services that regulate global climate and weather patterns.

Terrestrial ecosystems are predicted to receive unprecedented quantities of reactive N by the end of this century (Zak, Freedma, Upchurch, Steffens, & Kögel-Knabner, 2017). Recent studies on temperate and boreal forests have reported that elevated chronic N deposition significantly affects soil microbial abundance, community structure, and functional gene activity, and that the responses of soil microorganisms are strongly correlated with changes in ecosystem

functions including SOC cycling, both directly and indirectly, through multiple plant–soil–microbe interactions (Boot, Hall, Deneff, & Baron, 2016; Carrara et al., 2018; Eisenlord et al., 2013; Garcia-Palacios et al., 2015; Treseder, 2008; Zhang, Chen, & Ruan, 2018). NPP is stimulated by N deposition until N saturation is reached; increasing above- and belowground litter inputs (de Vries, Du, & Butterbach-Bahl, 2014). N fertilization may reduce C allocation belowground by modifying root physiology and exudation, and retarding rhizosphere priming effects (Dungait, Hopkins, Gregory, & Whitmore, 2012; Janssens et al., 2010; Kuzyakov, 2002; Zhu et al., 2014). Shifts in fungal community composition and functions were reported as a dominant mechanism driving soil C gains following N addition to temperate forest soils (Hassett, Zak, Blackwood, & Pregitzer, 2009; Hesse et al., 2015; Kellner, Luis, Schlitt, & Buscot, 2009). Decomposition rates are reduced by the effect of N fertilization on fungal activity as the genes encoding lignocellulolytic enzymes are downregulated (Chen et al., 2018; Hesse et al., 2015), or by inhibiting fungal growth (Treseder, 2008; Waldrop, Zak, Blackwood, Curtis, & Tilman, 2006). However, how anthropogenic N deposition affects soil microbial community structure and metabolic potential in tropical forests, and the subsequent impact on SOC accumulation, remains an open question (Cusack et al., 2016; Janssens et al., 2010; Pajares & Bohannan, 2016). Recent studies reported that N addition increased bacterial biomass, but decreased fungal/bacterial ratios in (sub)tropical forest soils, accompanied by detectable increases in complex organic SOC compounds (Cusack, Silver, Torn, Burton, & Firestone, 2011; Zhou, Wang, Zheng, Jiang, & Luo, 2017), although the opposite effect has also been observed (Li et al., 2015; Wang, Liu, & Bai, 2018).

The relatively N-rich status of tropical forests, compared to temperate/boreal forests, may moderate response intensity to N augmentation through atmospheric deposition (Cleveland et al., 2011; Hedin, Brookshire, Menge, & Barron, 2009). N dissimilative process rates are usually pronounced in N-rich ecosystems (Levy-Booth, Prescott, & Grayston, 2014), causing higher N₂O losses by denitrification and NO₃⁻ leaching from tropical soils compared to temperate forest soils (Vitousek & Matson, 1988; Zhang, Yu, Zhu, & Cai, 2014). Increasing N availability in tropical forests inhibits biological

N fixation (Cusack, Silver, & McDowell, 2009), alters nitrification (Han, Shen, Zhang, & Müller, 2018), and increases N_2O emissions (Liu & Greaver, 2009), which increase greenhouse gas (GHG) warming potential. Furthermore, chronic N deposition causes soil acidification with consequences for the availability and leaching of P, base cations (Mg^{2+} , K^+ , and Ca^{2+}), and micronutrients (Mo and Zn) that are essential for plants and microorganisms (Cusack et al., 2016; Lu et al., 2015; Zamanian, Zarebanadkouki, & Kuzyakov, 2018). Soil acidification can cause direct effects on plant and microbial community composition, and indirectly through the release of Al^{3+} , which has a broad-spectrum toxicity for plants (Kaspari et al., 2017). Wang et al. (2009) reported significantly increased Al^{3+} concentrations under high N addition soon after experimental treatment was started in a tropical forest. Soil acidification may reduce P transport across cell membranes, exacerbating the existing P limitation in weathered tropical forest soils (Kaspari et al., 2017; Li et al., 2016).

Atmospheric N deposited on "pristine" ecosystems, including old-growth tropical forests, may impact ecological functioning in remote locations, where monitoring is sparse and the consequences on biogeochemical cycling are difficult to determine (Holtgrieve et al., 2011). China has become the largest N creator and emitter globally (Liu, Zhang, Han, et al., 2013). Modeled predictions of N deposition are for 105 Tg N/year across Asia by 2030 (Mo et al., 2008; Zheng et al., 2002), and deposition rates of up to 73 kg N ha^{-1} year $^{-1}$ have been already recorded in tropical old-growth forests in southern China (Liu et al., 2011). While experimental data on the responses of temperate and boreal forests to elevated N deposition are relatively extensive, similar investigations in Chinese tropical forest ecosystems are sparse, and of a much shorter timescale, with the most well-established starting in the 2000s (Liu et al., 2011). However, tropical forest responses to long-term (decadal) atmospheric N fertilization, such as that currently experienced by natural ecosystems in China, and the effects on the functional ecology of such "pristine" tropical forests are likely to be profound. A recent meta-analysis by de Vries et al. (2014) on ecosystem N retention and C:N responses in forest ecosystem components (canopy, shoots and roots, and soil) suggests that the biological responses to increasing N deposition in tropical forests are different to those of boreal and temperate forests. However, we know very little about the functional ecology of the soil microorganisms that ultimately control the capacity of tropical forests to moderate ecosystem function in response to N deposition, and thereby act as a net SOC sink or nutrient cycling.

We exploited an existing 13-year N deposition field experiment in a monsoonal evergreen broadleaf old-growth forest in southern China. Previous research in the region has provided evidence that old-growth forests accumulated SOC over a 24-year period from 1979 to 2003 (Zhou et al., 2006). Previously, we reported that N addition increased rates of plant transpiration (Lu et al., 2018) and N_2O emissions (Zhang, Mo, Zhou, et al., 2008), but reduced litter decomposition (Fang, Mo, Peng, Li, & Wang, 2007) and soil respiration (Mo et al., 2008). We also found that N addition aggravated soil acidification and P supply, but increased Al mobility (Lu et al., 2015). However, the extent to which shifts in the microbial functional

community underpin these changes was not investigated. Here, we hypothesized that (a) long-term N additions alter microbial community composition by increasing soil acidification; and (b) the effects of N addition on the microbial community composition would translate into changes in microbial function that help explain the observed effects on soil C, N, and P cycling. To test these hypotheses, we used a microarray-based tool (GeoChip 5.0) to profile microbial functional potentials, specifically targeting a wide range of functional genes associated with C, N, and P cycling in soils. This method can be used as a specific and sensitive tool to study microbial functional potentials using correlations between microbial communities and ecosystem processes (He et al., 2010; Yang et al., 2013; Zhou et al., 2012).

2 | MATERIALS AND METHODS

2.1 | Field experiment and sampling

An N addition experiment was established in 2003 in a tropical old-growth monsoon evergreen broadleaf forest at the Dinghushan Biosphere Reserve (DHSBR), an UNESCO/MAB site located in the middle of Guangdong Province in southern China (23°10'N and 112°10'E; 250–300 m a.s.l.). The region has a monsoonal humid tropical climate with a mean annual temperature of 21°C (range 13°C in January to 28°C in July), and 1,930 mm of precipitation falling in a distinct seasonal pattern; 75% of the precipitation falls from March to August, and 6% from December to February. The soil is lateritic red earth formed from sandstone (Oxisol).

The experimental design is described in Mo et al. (2008). In brief, 12 plots of four treatments with three replicates were laid out in a completely randomized block design. Each plot was 20 m × 10 m, and was bounded by a 10-m wide buffer strip. N was added as ammonium nitrate (NH_4NO_3) solution at four N levels: Control (CK; zero N), Low-N (LN; 50 kg N ha^{-1} year $^{-1}$), Medium-N (MN; 100 kg N ha^{-1} year $^{-1}$), and High-N (HN; 150 kg N ha^{-1} year $^{-1}$). N was applied below the canopy on 12 occasions at monthly intervals each year at equal rates using a backpack sprayer. The control soil was treated with an equal volume (20 L) of deionized water.

The experimental plots were sampled in October 2015. Prior to soil sampling, CO_2 , N_2O , and CH_4 fluxes were collected from each plot from two static chambers that were inserted 5 cm into the soil (Zhang, Mo, Zhou, et al., 2008). Gas samples were taken using 100 ml plastic syringes at 0, 10, 20, and 30 min after chamber closure, and were analyzed in the laboratory within 24 hr.

Soil samples were taken using a soil corer (0–10 cm deep, 2 cm inner diameter) from 10 random points across each plot, and mixed to yield one composite sample per plot. The litter layer was carefully removed before sampling. The soil samples were stored in airtight polypropylene bags and placed in a cool box at 4°C during transportation to the laboratory. Litter, roots, and stones were carefully removed by hand, and the soil was divided into several subsamples. Subsamples for dissolved organic carbon (DOC), ammonium N (NH_4^+ -N), nitrate N (NO_3^- -N) and available phosphorus concentration analyses were stored at 4°C for no longer than 1 week. Subsamples for pH, SOC,

total nitrogen (TN), total phosphorus (TP), and pH analyses were air dried. Subsamples for microbial community composition and functional gene (GeoChip) analysis were stored at -80°C .

2.2 | Greenhouse gas and soil analyses

Greenhouse gas (CO_2 , N_2O , and CH_4) concentrations of the air sampled in each treatment were measured by gas chromatography (Agilent 4890D, Agilent Co.). Soil pH was measured with a pH meter after shaking the soil in deionized water (1:2.5 w/v) suspensions for 30 min. The SOC and TN content were determined by combustion using a Vario EL III Elemental Analyzer (Elementar). Ammonium N ($\text{NH}_4^+\text{-N}$) and $\text{NO}_3^-\text{-N}$ concentrations were determined using an autoanalyzer (TRAACS-2000, BRAN+LUEBBE) following 0.01 M KCL (1:10 w/v) extraction for 30 min. Total P and available P concentrations were measured using the ammonium molybdate method after $\text{H}_2\text{SO}_4\text{-H}_2\text{O}_2\text{-HF}$ digestion. The DOC concentration was measured following the method used by Jones and Willett (2006). The field-moist soil samples (equivalent to 15 g oven-dried soil) were extracted with 60 ml of 0.05 mol/L K_2SO_4 (soil to solution ratio 1:4) for 1 hr. The extract was then passed through a 0.45-mm membrane filter and analyzed for DOC using a Multi 3100 N/C TOC analyzer (Analytik). Microbial community composition was analyzed by phospholipid fatty acid (PLFA) composition according to Frostegård, Tunlid, and Bååth (1991). Changes in microbial community composition were presented as molar percentages (mole %) of the PLFA biomarkers for bacteria or fungi extracted from the soil samples, and the ratio of biomarkers for fungi and bacteria (fungi:bacteria), as previously described by Bossio and Scow (1998) and Högberg, Högberg, and Myrold (2007).

2.3 | Analyses of microbial functional communities

Total DNA was extracted from 0.5 g of well-mixed soil using the PowerSoil kit (MoBio Laboratories) according to the manufacturer's instructions. DNA quality and quantity were measured using a NanoDrop ND-1000 Spectrophotometer (NanoDrop Technologies Inc.), and with PicoGreen[®] using a FLUOstar[®] Optima microplate reader (BMG Labtech), respectively. DNA hybridization was performed using GeoChip 5.0 according to Yang et al. (2013). Briefly, DNA samples were labeled with Cy-5 fluorescent dye using a random priming method, and purified with the QIA quick purification kit (Qiagen). The DNA was dried in a SpeedVac (ThermoSavant) at 45°C for 45 min. GeoChip hybridization was carried out at 42°C for 16 hr on a MAUI[®] hybridization station (BioMicro). After purification, GeoChips were scanned by a NimbleGen MS200 scanner (Roche) at 633 nm, using a laser power and photomultiplier tube gain of 100% and 75%, respectively.

2.4 | Statistical analyses

Raw GeoChip data were analyzed using a data analysis pipeline as described previously (He et al., 2010; Yang et al., 2013). The data were logarithmically transformed, and then divided by the mean value of

each slide. A minimum of two valid values of three biological replicates (samples from the same treatment), were required for each gene. Spots that were flagged, or with a signal to noise ratio less than 2.0, were considered poor in quality and removed from statistical analysis.

Detrended correspondence analysis (DCA) was used to assess changes in the overall microbial functional community structure (based on GeoChip data). Adonis was further performed to confirm significant changes in microbial functional community structures in any pair of samples. To determine the relative importance of soil and plant factors in shaping microbial functional community structure, a canonical correspondence analysis (CCA)-based variation partitioning analysis (VPA) was implemented (Ramette & Tiedje, 2007). DCA and VPA statistical analyses were performed in R v.3.2.1 with the vegan package. Matrices of the pairwise distance between functional microbial community structure (Bray-Curtis) and Euclidean distance of environmental variables were also constructed in R using the vegan package.

To further link microbial functional genes with environmental variables and biogeochemical processes, partial least squares path modeling (PLS-PM) and Pearson correlations were performed. PLS-PM is a data analysis method for variables that can be summarized by use of latent variables, and the fact that linear relationships exist between latent variables (Sanchez, 2013). Each latent variable included one or more manifest variables, for example, inorganic N including $\text{NH}_4^+\text{-N}$ and $\text{NO}_3^-\text{-N}$. Each manifest variable has the degree of relative contribution degree shown as an arrow between manifest variable and latent variable in the analysis of path diagram. Models with different structures are evaluated using the goodness of fit (GOF) statistics, a measure of their overall predictive power with $\text{GOF} > 0.7$ considered acceptable values. The models were constructed using the "inner plot" function of the R package. PLS-PM was performed based on the reduction of the full models (Kou et al., 2017). The environmental drivers selected in the model were the main predictors according to their contribution for variation based on Random Forest analysis (% of increase of MSE). Random Forest analysis was conducted in R v.3.2.1 using the random Forest package (Delgado-Baquerizo et al., 2016).

The significant differences of environmental variables and signal intensities for selected C, N, and P cycling functional genes in soil sampled from four N addition levels were determined by one-way ANOVA, followed by the least significant difference test using SAS V8 (SAS Inc. 1996). Statistical significance was determined at $p < .05$ for all analyses.

3 | RESULTS

3.1 | Long-term effects of N addition on greenhouse gas emissions and soil properties

Compared with the control treatment (CK), the effects of N addition on (CO_2 , N_2O , and CH_4) fluxes were significant under High-N and Medium-N treatments, but not for Low-N (Table 1). CO_2 emission was 33% less under High-N compared with control. As compared with control, CH_4 uptake decreased by 64% and 63% in Medium-N and High-N treatments, but N_2O emission increased by 113% under High-N. All measured

TABLE 1 Effect of 13 years nitrogen (N) fertilizer addition to soil in a tropical forest at different rates on carbon (C), nitrogen, and phosphorus (P) pools, and soil microbial composition and greenhouse gas emission

	Treatment (kg N ha ⁻¹ year ⁻¹)			
	CK 0 (control)	Low-N 50	Medium-N 100	High-N 150
Greenhouse gas emissions				
CO ₂ (mg CO ₂ m ⁻² hr ⁻¹)	91.8 ± 4.05 a	88.2 ± 2.27 ab	85.4 ± 1.40 ab	61.2 ± 0.86 b
CH ₄ (μg C m ⁻² hr ⁻¹)	-51.2 ± 7.66 b	-40.0 ± 3.93 b	-18.4 ± 3.88 a	-19.1 ± 4.06 a
N ₂ O (μg N m ⁻² hr ⁻¹)	29.2 ± 2.14 b	41.7 ± 5.16 ab	39.6 ± 9.28 ab	62.2 ± 9.48 a
Soil chemistry				
pH (H ₂ O)	3.9 ± 0.03 a	3.8 ± 0.05 ab	3.8 ± 0.02 b	3.7 ± 0.01 b
Soil organic carbon (g C ⁻¹ kg ⁻¹)	25.4 ± 3.44 b	29.6 ± 2.54 ab	30.2 ± 0.46 ab	31.9 ± 0.77 a
Dissolved organic carbon (mg C ⁻¹ g ⁻¹)	0.5 ± 0.04 a	0.6 ± 0.04 a	0.6 ± 0.04 a	0.6 ± 0.01 a
Total nitrogen (g N ⁻¹ kg ⁻¹)	1.8 ± 0.14 b	2.3 ± 0.28 ab	2.2 ± 0.07 ab	2.5 ± 0.08 a
Nitrate N (mg N ⁻¹ kg ⁻¹)	2.4 ± 0.27 c	9.6 ± 1.49 b	16.1 ± 2.83 a	17.7 ± 1.21 a
Ammonium N (mg N ⁻¹ kg ⁻¹)	4.0 ± 0.18 b	5.0 ± 0.53 ab	5.1 ± 0.72 ab	7.4 ± 1.66 a
Available phosphorus (mg P/kg)	0.5 ± 0.05 a	0.4 ± 0.08 ab	0.4 ± 0.02 b	0.4 ± 0.01 b
Total phosphorus (mg P/g)	0.2 ± 0.03 a	0.2 ± 0.01 a	0.2 ± 0.01 a	0.2 ± 0.01 a
Soil microbial composition (phospholipid fatty acids)				
Bacterial abundance (mole %)	52.4 ± 0.24 a	52.0 ± 0.31 a	51.4 ± 0.24 ab	50.0 ± 0.24 b
Fungal abundance (mole %)	10.0 ± 0.51 a	9.3 ± 0.16 a	9.9 ± 0.14 a	9.7 ± 0.39 a
Fungal:Bacterial ratio	0.19 ± 0.01 a	0.18 ± 0.00 a	0.19 ± 0.00 a	0.19 ± 0.01 a

Note: Values are presented as mean (n = 3) ± 1 SEM followed by lowercase letters that indicate significant difference among treatments ($p < .05$).

soil parameters in the High-N-treated soils aside from DOC and TP were significantly different from CK. SOC and TN contents and NH₄⁺ concentrations were higher under the High-N treatment. Nitrate (NO₃⁻) concentrations increased under Medium-N and High-N compared to Low-N and CK. Available P and pH were lower under Medium-N and High-N treatments than under Low-N. Bacterial relative abundance based on PLFA biomarker contents was lower under High-N treatment than under Low-N, but there was no significant difference in fungal PLFA biomarker concentrations or fungal:bacterial (F:B) ratios between treatments.

3.2 | Long-term effects of N addition on microbial functional community structure

The α diversity of the microbial functional community decreased with N application rate and was lowest under the High-N treatment (Figure S1; $p < .05$). N addition markedly changed soil microbial functional community structure, as indicated by DCA (Figure 1), which was confirmed by nonparametric multivariate statistical tests (Figure 1; $p < .001$). Further pairwise comparison revealed that microbial functional community structures under Medium-N and High-N treatments differed from CK (Table S1).

A partial CCA-based VPA evaluated the relative contributions of environmental factors to the microbial functional community structure (Figure S2). A total of 89.9% of the community variations could be explained by all measured variables. Soil and plant variables explained 72.6% and 11.8% of variations, respectively,




FIGURE 1 Detrended correspondence analysis of soil microbial functional community based on GeoChip data after analysis of tropical forest soils treated with different rates of N fertilizer for 13 years. Treatment definitions: CK (zero N addition; control); LN (Low-N, 50 kg N ha⁻¹ year⁻¹); MN (Medium-N, 100 kg N ha⁻¹ year⁻¹); HN (High-N, 150 kg N ha⁻¹ year⁻¹)

while their interactions explained 5.50% (Figure S2). Among soil variables, NO₃⁻-N and available P were the most important factors, contributing 14.7% and 16.2%, respectively, to variation in microbial functional community structure (Figure S2).

3.3 | Linking microbial functional genes to soil CO₂ emission

Key genes associated with C degradation were analyzed and related to soil CO₂ emissions. High N decreased the abundances of genes responsible for labile and recalcitrant C (starch, cellulose, and chitin) degradation ($p < .05$; Figure 2). For example, the relative abundances of *apu*, *npIT*, *amyX*, *cellobiase*, *acetylglucosaminidase*, and *chitinase* were the lowest under High-N ($p < .05$;

Figure S3). PLS path modeling explained 79% of the CO₂ emission variance, and provided the best fit to our data (GOF of 0.77; Figure 3a). Bacterial relative abundance showed the largest effect on CO₂ emission via direct (path coefficient = 0.34) and indirect effects (path coefficient = 0.62) on microbial functional C degradation genes (Figure 3a). There were corresponding strong, positive correlations between bacterial relative abundance and the abundances of genes involved in starch, hemicelluloses, cellulose, and chitin degradation, but no similar relationship was




FIGURE 2 Normalized signal intensity of detected genes indicating the degradation of different organic compounds (a–e) in tropical soils treated with different rates of N fertilizer addition (CK, LN, MN and HN; see Figure 1 for treatment definitions) for 13 years. Signal intensities were summed and normalized by the probe number for each substance. Different lowercase letters indicate significant differences among treatments; error bars indicate standard error of the mean ($n = 3$)




FIGURE 3 Partial least squares path analysis for greenhouse gas (GHG) fluxes—(a) CO₂, (b) N₂O, and (c) CH₄—from a tropical forest soil, showing the relationships between selected biogeochemical processes, microbial community composition, and functional gene abundances. GOF, goodness of fit. GHG pools are shown in blue, microbial functional genes in black, and environmental variables in yellow green. The blue arrows are the direct effect of environmental and microbial variables on GHG fluxes, and the black arrows indicate the indirect path. The numbers listed within arrows are the standardized path coefficient [Colour figure can be viewed at wileyonlinelibrary.com]

observed with fungal relative abundance (Figure 4). The positive direct effect of C degradation genes (path coefficient = 0.57) on CO₂ emission was the greatest (Figure 3a). Among C degradation genes, strong positive correlations between cellulose and chitin degradation gene abundances and CO₂ production were observed (Figure S4; $p < .05$). This finding was consistent with PLS path modeling that showed large loading factors (Figure 3a).

3.4 | Linking microbial functional genes to soil N₂O emission

With increasing N levels, the abundance of *gdh* gene (glutamate dehydrogenase) decreased, while that of *ureC* (urease) increased (Figure 5; $p < .05$), suggesting a shift in microbial functional potential toward N mineralization. Both genes associated with nitrification (*amoA* and *hao*) decreased under Medium-N and High-N treatments. High-N increased denitrification genes including *narG* (membrane-bound nitrate reductase), *norB* (nitric oxide reductase), and *nirK* (copper-containing nitrite reductase). Correspondingly, NO₃⁻-N concentrations were related to the increased abundance of genes encoding *NiR* and *nirA* (assimilatory nitrate reduction). The abundance of the N fixation gene (*nifH*) was the lowest under High-N treatment.

PLS path modeling explained 89% of the N₂O emission variance (Figure 3b). There was a direct positive effect of denitrification genes on N₂O emission in the PLS path model that was supported by the Pearson correlations (Figure S5). This suggested that increased denitrification was the driving force for N₂O production after N amendment. Among soil variables, the positive direct effect of inorganic N concentrations on N₂O emission was the greatest, while pH was related to the largest negative indirect effect (Figure 3b).

3.5 | Linking microbial functional genes to soil CH₄ uptake

High-N increased the abundance of the gene encoding *mcrA* (methyl coenzyme M reductase), a key enzyme in methanogenesis (Figure 6a; $p < .05$). In contrast, there was a higher abundance of genes encoding *mmoX* (involved in methane oxidation) under the Low-N treatment. PLS path modeling accounted for 81% of the variation in CH₄ uptake (Figure 3c). There were strong positive direct effects of *mcrA* (methanogenesis) on CH₄ uptake in PLS path modeling (Figure 3c). In general, the direct effect of DOC (path coefficient = 0.48) and pH (path coefficient = -0.50) on CH₄ flux was greater than those of SOC (path coefficient = -0.28) and inorganic N concentrations (path coefficient = -0.13; Figure 3c).

3.6 | Changes in key genes related to P cycling

The relative abundance of the phytic acid hydrolysis gene (encoding phytase) decreased under Medium-N and High-N additions (Figure 6b; $p < .05$). Most genes encoding phytase were derived from Proteobacteria. The decrease in microbial P utilization genes after N amendment suggested that N amendments exacerbate soil P deficiency. This was supported by positive relationships between phytase abundance, and available and total P contents ($p < .05$; Table S2). We found positive correlations between phytase abundance and soil pH, but negative correlations with inorganic N concentrations ($p < .05$; Table S2).




FIGURE 4 Pearson correlations between abundances of C degradation genes indicating potential decomposition of organic C compounds (a–l) in a tropical forest soil and relative abundances of bacteria and fungi




FIGURE 5 The normalized average signal intensity of detected functional genes for N cycling (a–f) after N fertilizer addition for 13 years at different rates (CK, LN, MN, HN; see Figure 1 for treatment definitions). Signal intensities were averaged and normalized by the probe number for each gene. Different lowercase letters indicate significant differences among treatments; error bars indicate standard error of the mean ($n = 3$) [Colour figure can be viewed at wileyonlinelibrary.com]




FIGURE 6 The normalized average signal intensity of detected genes indicating CH₄ (a) and P (b) cycling genes in tropical forest soils treated with different rates of N fertilizer for 13 years (CK, LN, MN, HN; see Figure 1 for treatment definitions). Signal intensities were averaged and normalized by the probe number for each gene. Different lowercase letters indicate significant differences among treatments; error bars indicate standard error of the mean ($n = 3$) [Colour figure can be viewed at wileyonlinelibrary.com]

4 | DISCUSSION

Long-term (13-year) NH₄NO₃ addition in a tropical old-growth forest altered microbial composition, predominantly by decreasing the relative abundance of bacteria (Table 1). Furthermore, microbial functional mechanisms encoded by C, N, and P cycling genes were altered by N addition at high rate, and explained the observed effects on soil C, N, and P cycling at the experimental site (summarized in Figure 7). Therefore, our hypotheses regarding the significant changes in microbial composition and functions after N deposition at rates higher than 100 kg N ha⁻¹ year⁻¹ were confirmed. The potential impacts of these findings are discussed in the context of reported phenomena pertinent to the climate change mitigation potential of tropical old-growth forests under chronic N addition, that is, SOC accumulation and increased N loss via denitrification.

4.1 | SOC accumulation in old-growth tropical forests under N addition

The positive priming of SOM and increased soil respiration rates caused by increases in NPP and organic matter input after atmospheric CO₂ enrichment of tropical forests reported by Sayer, Heard, Grant, Marthews, and Tanner (2011) is apparently contradicted by the observed increase in SOC accumulation after N addition in old-growth tropical forest under High-N in this study. No changes in plant growth and litter input were observed at the DHSBR experimental site (Mo et al., 2008). Furthermore, Mo et al. (2008) showed that High-N addition inhibited fine root growth at the same experimental site. Declining root biomass is a key response to N saturation,




FIGURE 7 A conceptual diagram illustrating the positive (+) or negative (-) impact of high N addition on the potential activity of microbial functional genes that control C and N cycling in tropical forest soils after N addition for 13 years. Greenhouse gas pools are shown in green, substrate pools in yellow, microbial functional genes in orange, and biological processes in blue

and has also been reported by Magill et al. (2004) in the chronic N amendment in the Harvard Forest, USA, where similar rates ($50\text{--}150\text{ kg N ha}^{-1}\text{ year}^{-1}$) were applied to hardwood and pine forest soils. Therefore, changes in aboveground NPP or belowground NPP are unlikely to provide a route for increased SOC in the DHSBR tropical forest under increased N.

Undisturbed forest soils have large F:B ratios (e.g. 0.3–0.5; Frostegård & Bååth, 1996), and large chronic inorganic N input reduces fungal biomass and inhibits lignases (Frey, Knorr, Parrent, & Simpson, 2004; Hassett et al., 2009). Zak et al. (2017) recently proposed that reduced decomposition in temperate forests is caused by a shift from basidiomycete fungal activity to bacteria and ascomycete fungi that only can partially oxidize polyphenolic structures in soil. In our study, however, there were no differences between the fungal relative abundance, the F:B ratio (approximately 0.2; Table 1), or the gene abundances of ligninolytic enzymes between N addition levels (Figure 2). However, the nonexperimental N addition (i.e. annual atmospheric addition of $40\text{--}50\text{ kg N ha}^{-1}\text{ year}^{-1}$ recorded at the site) had already reduced the fungal population in the control plots preceding experimental N addition. Indeed, a moderate increase of N addition ($15\text{ kg N ha}^{-1}\text{ year}^{-1}$) for 5 years from 2007 to 2011 in DHSBR soils increased fungal biomass and F:B ratios, which were also measured using PLFA analysis (Liu,

Zhang, Han, et al., 2013). This may suggest that the larger N addition in our study exceeded a biological threshold controlling the activity of soil fungi. Similarly, Frey et al. (2004) reported that active fungal biomass was 27%–69% reduced in the Harvard Forest experiment where background atmospheric N input was relatively low ($8\text{ kg N ha}^{-1}\text{ year}^{-1}$) but bacterial biomass was unchanged, resulting in lower F:B ratios. Högborg et al. (2007) also determined significant decreases in the mole % of fungal biomass and F:B ratio (similarly based on PLFA analysis of 18:2ω6) in a northern European boreal forest after long-term (20+ years) N addition ($34\text{--}108\text{ kg N ha}^{-1}\text{ year}^{-1}$), where the background atmospheric N deposition was minimal ($3\text{ kg N ha}^{-1}\text{ year}^{-1}$). This suggests that chronic N addition has a negative effect on fungal survival in forest soil. The effect of a reduction in rhizosphere exudation as a key mechanism for the substantial (45%) decrease in ectomycorrhizal fungi biomass was suggested by comparison with experimental tree girdling, providing an explanation for the decrease in microbial heterotrophic respiration previously observed in a Chinese pine forest (Fan et al., 2014; Wang et al., 2016). Therefore, increased root and active fungal biomass, as observed by Clemmensun et al. (2013) for N-limited boreal forest ecosystems, are unlikely to be the primary drivers of SOC accumulation in DHSBR tropical forest soils under chronic N addition.

High-N addition decreased CO₂ emission by 33% in this study (Table 1), concurring with previous studies that reported that N additions decreased soil respiration in recent studies of tropical forest soils (Fan et al., 2014; Wang et al., 2016), although by comparison, Feng et al. (2017) reported no change. We observed that bacterial relative abundance was decreased under High-N addition (Table 1). Wang et al. (2009) explored the initial effects of N addition in the early stages of the experiment and found that soil microbial biomass C (extracted by chloroform fumigation) was generally decreased in N addition plots, but significantly decreased in the High-N plots after 2 and 4 years. These observations are contrary to the observations of similar experiments in temperate and boreal forests in general. Furthermore, bacterial relative abundance had the largest total effect on total soil respiration (CO₂ emission; Figure 3a; $p < .05$). A decrease in microbial abundance provides a straightforward explanation for the measured decrease in CO₂ emissions from tropical forests under High-N addition, as previously proposed by Wei, Jiangming, Yunting, Xiankai, and Hui (2008). Direct and indirect effects of each latent variable on target variables were identified using PLSPM analysis. The processes underlying the measured CO₂ emission in the field include soil organic matter, roots, and microorganisms derived (Kuzyakov, 2006). Thus, the indirect effect of bacteria on CO₂ may be through changes in the production of C degradation enzymes, and the direct effect of bacteria on CO₂ may derive from changes in microbial respiration, in response to N addition. Significant relationships between CO₂ emission and the relative abundances of cellulose and chitin degradation genes were observed, respectively (Figure S4; $p < .05$). This indicates that N addition inhibited the production of C degradation enzymes by the soil bacterial biomass, which may drive the accumulation of intact organic compounds in the soil, thereby increasing SOC under High-N conditions. This is further evidence supporting the hypothesis that SOM turnover is controlled by the access of microorganisms and their enzymes, regardless of assumed chemical recalcitrance (Dungait et al., 2012).

Tropical forest soils often have a limited capacity to buffer acidification, so N addition can lead to rapid soil acidification (Lu et al., 2015; Matson, McDowell, Townsend, & Vitousek, 1999; Zamanian et al., 2018), negatively affecting bacterial growth by an increase in toxic free Al³⁺ (Rousk, Brookes, & Baath, 2009), but also the loss of other nutrients including P and cations (Mg²⁺, Ca²⁺) from the soil by leaching. The available P concentrations in High-N treatment reveal a decrease as compared to CK (Table 1). Magill et al. (2004) observed Ca²⁺, Mg²⁺ and K⁺ leaching from the O horizon under N addition, and subsequent declines in foliar Mg:N and Ca:Al in the Harvard Forest, suggesting a positive feedback to decrease biological availability with time. Our results concur with previous studies suggesting bacteria are less tolerant of acidic environments than fungi (Li et al., 2015; Wang et al., 2018). Overall, our study demonstrates that long-term High-N addition in a tropical forest promotes net C gain by altering microbial community composition and inhibiting organic C degradation functional potential, leading to an accumulation

of undegraded organic compounds. However, this C sequestration potential needs to be considered against the changes observed for other aspects of soil biogeochemistry, for example, macro- and micronutrient cycling, and the wider impacts on ecosystem functions including biodiversity.

According to the in situ measurements of CH₄ fluxes, the soils under all treatments acted as net sinks of CH₄, but this was significantly less in the plots treated with high or medium rates of N fertilizer application (−18.4 and −19.1 μg C m^{−2} hr^{−1}, respectively) compared to low N addition or the control (−40.0 and −51.2 μg C m^{−2} hr^{−1}, respectively; Table 1). This indicates that the balance between methane methanotrophy and methanogenesis in the tropical forest soils in this study had been affected by N addition. The global meta-analysis performed by Liu and Greaver (2009) showed that N addition increased CH₄ emission by 97% and reduced CH₄ uptake by 38%. Schimel (2000) had previously described that high NH₄⁺ concentrations could inhibit CH₄ oxidation because of competition for methane monooxygenase resulting in increased CH₄ emissions. Methanotrophy indicated by *pmoA* and *mmoX* expression was not significantly different from the control aside from *mmoX* under LN, but *mcrA* expression (methanogenesis) was increased under HN (Figure 6). Additionally, in this study, the strong positive direct effect of *mcrA* on CH₄ uptake (Figure 3c) suggests that the decrease of CH₄ uptake may largely be due to increased methanogenesis. The abundance of the *mcrA* gene positively correlates with CH₄ production potential (Ma, Conrad, & Lu, 2012). Accelerated methanogenesis could be partially attributed to increased C availability; we found that labile C (DOC) had a strong direct effect on the *mcrA* gene and CH₄ flux (Figure 3c). This agrees with previous reports that labile C pool fuels the activity of methanogenic populations, resulting in increased CH₄ emissions (Dorodnikov, Knorr, Kuzyakov, & Wilmking, 2011; Lu, Lueders, Friedrich, & Conrad, 2005; Zheng et al., 2017).

4.2 | Increased potential N loss in old-growth tropical forests under chronic N addition

Soil C and N cycles are closely linked, and often isometric in their stoichiometric expression (Beniston et al., 2015). Therefore, the observation that SOC is accumulating under N addition in tropical forest soils infers that organic N will contribute to increased N mineralization, as attested by the increased relative abundances of *ureC* genes that control ammonification under High-N conditions (Figure 5), and the positive correlations between ammonification gene abundance and DOC ($r = .55$, $p < .05$). However, *gdh* expression decreased under High-N conditions, suggesting the two genes encoding autotrophic ammonia-oxidizing metabolic traits may be expressed in organisms that respond differentially to N addition. Ammonia-oxidizing archaea dominate nitrification (Prosser & Nicol, 2012) and *ureC* expression in acid soils (Lu & Jia, 2013), suggesting a switch from bacterial to increasing archaeal functional potential as bacterial abundance decreased under High-N conditions.

The N_2O flux from soil mainly originates from nitrification or denitrification (Levy-Boothet et al., 2014). The large increase in N_2O emissions (113%; Table 1) under High-N conditions coincided with increased soil acidity, the metabolic potential for NO_3^- -N reduction in this study (Figure 3b), and an increased relative abundance of genes involved in denitrification (*narG* and *norB*) and assimilatory N reduction (*NiR* and *nirA*; Figure 3b, Figures S5 and S7). The strong direct positive effect of denitrification genes, but not nitrification genes, on N_2O emission (path coefficient = 0.67; Figure 3b) suggests that denitrification was a driving force of N_2O emission in this acidic tropical forest soil. Our result agreed with previous studies that N addition can increase N_2O emissions and is correlated with genes involved in the denitrification pathway (Corre, Veldkamp, Arnold, & Wright, 2010; Han et al., 2018). Among denitrification genes, we determined significant contributions from *narG* and *norB* genes to N_2O emissions (Figure 3b; Figure S5). Previous studies found that the *nosZ* gene was positively correlated with soil NO and N_2O production in acidic forests (Lammel, Feigl, Cerri, & Nusslein, 2015; Yu et al., 2014), but the abundance of *nosZ* was not significantly different from the control in our study. Relative abundances of *nirK* were reported to increase under enhanced NO_3^- -N concentrations (Morales, Cosart, & Holben, 2010) and low pH (Rütting, Huygens, Boeckx, Staelens, & Klemmedtsson, 2013). Nitrogenous gas emissions (N_2O/N_2 emission ratios) can be influenced by measurable soil properties in tropical soils including parent materials, pH, soil moisture, and redox potential (Kang, Mulder, Duan, & Dorsch, 2017; Liu, Morkved, Frostegård, & Bakken, 2010; Rütting et al., 2013; Stone, Kan, & Plante, 2015; Zhang, Cai, Cheng, & Zhu, 2009). Therefore, the variations between soil N processes and gene expression response to N additions are complex and barely predictable (Cheng et al., 2019; Levy-Boothet et al., 2014). Indeed, in our experiment, *nirS* decreased, again suggesting that ubiquitous soil processes including denitrification are performed by a range of microorganisms that respond differently to stresses including pH change. The decreased abundance of nitrification genes that coincided with the increased abundances of denitrification genes and N_2O emissions indicate a potentially high N loss, likely counteracting net N accumulation in this study. Most significantly, considering the high warming potential of N_2O , the stimulated N_2O emissions potentially offset the perceived beneficial effect of N addition on soil C accumulation in tropical old-growth forest soils, and suggest that the effects of chronic N deposition on remote ecosystems must be considered in models predicting feedbacks to climate change.

In conclusion, this study revealed profound interrelationships between the response of soil microbial functional potentials and soil C, N, and P cycling to chronic N addition in "pristine" tropical forests. Changes in bacterial relative abundance significantly affected CO_2 emissions, mainly via indirect effects on microbial functional C degradation genes leading to SOC accumulation. We revealed the metabolic potential for the increased expression of microbial N functional genes driving increased N_2O fluxes from these ecosystems.

Limitations on P availability caused by soil acidification through reduced *phytase* expression were also identified. This new understanding of the effect of human-induced atmospheric N deposition must be factored into Earth system models considering the GHG sink capacity of tropical forests, and the effect of air pollution from agriculture and industry on the biogeochemical processes and biodiversity of tropical ecosystems. Further studies are necessary to investigate the effect of climatic variation at a range of timescales (seasonal and decadal) to investigate the impact of N addition on the dynamics of microbial communities and functions, and their roles in mediating soil biogeochemistry, under a more extensive range of environmental conditions experienced by the soil microbial community.

ACKNOWLEDGEMENTS

This study was supported by the National Natural Science Foundation of China (grant nos. 31770560, 41571130041 and 41731176), the National Key R&D Program of China (grant no. 2017YFA0604803), and the Youth Innovation Research Team Project (grant no. LENOM2016Q0004). Yakov Kuzyakov is very thankful for the Changjiang Scholar Award.

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

ORCID

Yunfeng Yang  <https://orcid.org/0000-0001-8274-6196>

Jiangming Mo  <https://orcid.org/0000-0002-1437-8018>

Guirui Yu  <https://orcid.org/0000-0002-1859-8966>

Yakov Kuzyakov  <https://orcid.org/0000-0002-9863-8461>

REFERENCES

- Baccini, A., Walker, W., Carvalho, L., Farina, M., Sulla-Menashe, D., & Houghton, R. A. (2017). Tropical forests are a net carbon source based on aboveground measurements of gain and loss. *Science*, 358, 230–233. <https://doi.org/10.1126/science.aam5962>
- Beniston, J. W., Shipitalo, M. J., Lal, R., Dayton, E. A., Hopkins, D. W., Jones, F., ... Dungait, J. A. J. (2015). Carbon and macronutrient losses during accelerated erosion under different tillage and residue management. *European Journal of Soil Science*, 66, 218–225. <https://doi.org/10.1111/ejss.12205>
- Boot, C. M., Hall, E. K., Deneff, K., & Baron, J. S. (2016). Long-term reactive nitrogen loading alters soil carbon and microbial community properties in a subalpine forest ecosystem. *Soil Biology & Biochemistry*, 92, 211–220. <https://doi.org/10.1016/j.soilbio.2015.10.002>
- Bossio, D. A., & Scow, K. M. (1998). Impacts of carbon and flooding on soil microbial communities: Phospholipid fatty acid profiles and substrate utilization patterns. *Microbial Ecology*, 35(3), 265–278. <https://doi.org/10.1007/s002489900082>
- Camenzind, T., Hattenschwiler, S., Treseder, K. K., Lehmann, A., & Rillig, M. C. (2018). Nutrient limitation of soil microbial processes in tropical forests. *Ecology Monographs*, 88, 4–21. <https://doi.org/10.1002/ecm.1279>

- Carrara, J. E., Walter, C. A., Hawkins, J. S., Peterjohn, W. T., Averill, C., & Brzostek, E. R. (2018). Interactions among plants, bacteria, and fungi reduce extracellular enzyme activities under long-term N fertilization. *Global Change Biology*, 24, 2721–2734. <https://doi.org/10.1111/gcb.14081>
- Chen, J., Luo, Y. Q., Van Groenigen, K. J., Hungate, B. A., Cao, J. J., Zhou, X. H., & Wang, R. W. (2018). A keystone microbial enzyme for nitrogen control of soil carbon storage. *Science Advance*, 4, eaa1689. <https://doi.org/10.1126/sciadv.aaq1689>
- Cheng, Y., Wang, J., Chang, S. X., Cai, Z. C., Muller, C., & Zhang, J. B. (2019). Nitrogen deposition affects both net and gross soil nitrogen transformations in forest ecosystems: A review. *Environmental Pollution*, 244, 608–616. <https://doi.org/10.1016/j.envpol.2018.10.054>
- Clemmensen, K. E., Bahr, A., Ovaskainen, O., Dahlberg, A., Ekblad, A., Wallander, H., ... Lindahl, B. D. (2013). Roots and associated fungi drive long-term carbon sequestration in boreal forest. *Science*, 339, 1615–1618. <https://doi.org/10.1126/science.1231923>
- Cleveland, C. C., Townsend, A. R., Taylor, P., Alvarez-Clare, S., Bustamante, M. M. C., Chuyong, G., ... Wieder, W. R. (2011). Relationships among net primary productivity, nutrients and climate in tropical rain forest: A pan-tropical analysis. *Ecology Letters*, 14, 939–947. <https://doi.org/10.1111/j.1461-0248.2011.01658.x>
- Corre, M. D., Veldkamp, E., Arnold, J., & Wright, S. J. (2010). Impact of elevated N input on soil N cycling and losses in old-growth lowland and montane forests in Panama. *Ecology*, 91, 1715–1729. <https://doi.org/10.1890/09-0274.1>
- Cusack, D. F., Karpman, J., Ashdown, D., Cao, Q., Ciochina, M., Halterman, S., ... Neupane, A. (2016). Global change effects on humid tropical forests: Evidence for biogeochemical and biodiversity shifts at an ecosystem scale. *Reviews of Geophysics*, 54, 523–610. <https://doi.org/10.1002/2015RG000510>
- Cusack, D. F., Silver, W., & McDowell, W. H. (2009). Biological nitrogen fixation in two tropical forests: Ecosystem-level patterns and effects of nitrogen fertilization. *Ecosystems*, 12, 1299–1315. <https://doi.org/10.1007/s10021-009-9290-0>
- Cusack, D. F., Silver, W. L., Torn, M. S., Burton, S. D., & Firestone, M. K. (2011). Changes in microbial community characteristics and soil organic matter with nitrogen additions in two tropical forests. *Ecology*, 92, 621–632. <https://doi.org/10.1890/10-0459.1>
- De Vries, W., Du, E. Z., & Butterbach-Bahl, K. (2014). Short and long-term impacts of nitrogen deposition on carbon sequestration by forest ecosystems. *Current Opinion in Environmental Sustainability*, 9–10, 90–104. <https://doi.org/10.1016/j.cosust.2014.09.001>
- Delgado-Baquerizo, M., Maestre, F. T., Reich, P. B., Jeffries, T. C., Gaitan, J. J., Encinar, D., ... Singh, B. K. (2016). Microbial diversity drives multifunctionality in terrestrial ecosystems. *Nature Communications*, 7. <https://doi.org/10.1038/ncomms10541>
- Dorodnikov, M., Knorr, K. H., Kuzyakov, Y., & Wilmking, M. (2011). Plant-mediated CH₄ transport and contribution of photosynthates to methanogenesis at a boreal mire: A ¹⁴C pulse-labeling study. *Biogeosciences*, 8, 2365–2375. <https://doi.org/10.5194/bg-8-2365-2011>
- Dungait, J. A. J., Hopkins, D. W., Gregory, A. S., & Whitmore, A. P. (2012). Soil organic matter turnover is governed by accessibility not recalcitrance. *Global Change Biology*, 18, 1781–1796. <https://doi.org/10.1111/j.1365-2486.2012.02665.x>
- Eisenlord, S. D., Freedman, Z., Zak, D. R., Xue, K., He, Z., & Zhou, J. (2013). Microbial mechanisms mediating increased soil C storage under elevated atmospheric N deposition. *Applied and Environmental Microbiology*, 79, 2847. <https://doi.org/10.1128/aem.00468-13>
- Fan, H. B., Wu, J. P., Liu, W. F., Yuan, Y. H., Huang, R. Z., Liao, Y. C., & Li, Y. Y. (2014). Nitrogen deposition promotes ecosystem carbon accumulation by reducing soil carbon emission in a subtropical forest. *Plant and Soil*, 379, 361–371. <https://doi.org/10.1007/s11104-014-2076-y>
- Fang, H., Mo, J., Peng, S., Li, Z., & Wang, H. (2007). Cumulative effects of nitrogen additions on litter decomposition in three tropical forests in southern China. *Plant and Soil*, 297, 233–242. <https://doi.org/10.1007/s11104-007-9339-9>
- Feng, J. G., Wang, J. S., Ding, L. B., Yao, P. P., Qiao, M. P., & Yao, S. C. (2017). Meta-analyses of the effects of major global change drivers on soil respiration across China. *Atmospheric Environment*, 150, 181–186. <https://doi.org/10.1016/j.atmosenv.2016.11.060>
- Frey, S. D., Knorr, M., Parrent, J. L., & Simpson, R. T. (2004). Chronic nitrogen enrichment affects the structure and function of the soil microbial community in temperate hardwood and pine forests. *Forest Ecology and Management*, 196, 159–171. <https://doi.org/10.1016/j.foreco.2004.03.018>
- Frostegård, A., & Bååth, E. (1996). The use of phospholipid fatty acid analysis to estimate bacterial and fungal biomass in soil. *Biology and Fertility of Soils*, 22, 59–65. <https://doi.org/10.1007/BF00384433>
- Frostegård, Å., Tunlid, A., & Bååth, E. (1991). Microbial biomass measured as total lipid phosphate in soils of different organic content. *Journal of Microbiological Methods*, 14, 151–163. [https://doi.org/10.1016/0167-7012\(91\)90018-L](https://doi.org/10.1016/0167-7012(91)90018-L)
- Galloway, J. N., Dentener, F. J., Capone, D. G., Boyer, E. W., Howarth, R. W., Seitzinger, S. P., ... Vorosmarty, C. J. (2004). Nitrogen cycles: Past, present, and future. *Biogeochemistry*, 70, 153–226. <https://doi.org/10.1007/s10533-004-0370-0>
- Garcia-Palacios, P., Vandegehuchte, M. L., Shaw, E. A., Dam, M., Post, K. H., Ramirez, K. S., ... Wall, D. H. (2015). Are there links between responses of soil microbes and ecosystem functioning to elevated CO₂, N deposition and warming? A global perspective. *Global Change Biology*, 21, 1590–1600. <https://doi.org/10.1111/gcb.12788>
- Han, X., Shen, W., Zhang, J., & Müller, C. (2018). Microbial adaptation to long-term N supply prevents large responses in N dynamics and N losses of a subtropical forest. *Science of the Total Environment*, 626, 1175–1187. <https://doi.org/10.1016/j.scitotenv.2018.01.132>
- Hassett, J. E., Zak, D. R., Blackwo, C. B., & Pregitzer, K. S. (2009). Are basidiomycete laccase gene abundance and composition related to reduced lignolytic activity under elevated atmospheric NO₃⁻ deposition in a northern hardwood forest? *Microbial Ecology*, 57, 728–739. <https://doi.org/10.1007/s00248-008-9440-5>
- He, Z., Deng, Y. E., Van Nostrand, J. D., Tu, Q. C., Xu, M. Y., Hemme, C. L., ... Zhou, J. Z. (2010). GeoChip 3.0 as a high-throughput tool for analyzing microbial community composition, structure and functional activity. *The ISME Journal*, 4, 1167–1179. <https://doi.org/10.1038/ismej.2010.46>
- Hedin, L. O., Brookshire, E. N. J., Menge, D. N. L., & Barron, A. R. (2009). The nitrogen paradox in tropical forest ecosystems. *Annual Review of Ecology Evolution and Systematics*, 40, 613–635. <https://doi.org/10.1146/annurev.ecolsys.37.091305.110246>
- Hesse, C. N., Mueller, R. C., Vuysich, M., Gallegos-Graves, L., Gleasner, C. D., Zak, D. R., & Kuskel, C. R. (2015). Forest floor community metatranscriptomes identify fungal and bacterial responses to N deposition in two maple forests. *Frontiers in Microbiology*, 6, 337. <https://doi.org/10.3389/fmicb.2015.00337>
- Högberg, M. N., Högberg, P., & Myrold, D. D. (2007). Is microbial community composition in boreal forest soils determined by pH, C-to-N ratio, the trees, or all three? *Oecologia*, 150, 590–601. <https://doi.org/10.1007/s00442-006-0562-5>
- Holtgrieve, G. W., Schindler, D. E., Hobbs, W. O., Leavitt, P. R., Ward, E. J., Bunting, L., ... Wolfe, A. P. (2011). A coherent signature of anthropogenic nitrogen deposition to remote watersheds of the northern hemisphere. *Science*, 334, 1545–1548. <https://doi.org/10.1126/science.1212267>
- Janssens, I. A., Dieleman, W., Luysaert, S., Subke, J. A., Reichstein, M., Ceulemans, R., ... Law, B. E. (2010). Reduction of forest soil respiration in response to nitrogen deposition. *Nature Geoscience*, 3, 315–322. <https://doi.org/10.1038/ngeo844>

- Jones, D. L., & Willett, V. B. (2006). Experimental evaluation of methods to quantify dissolved organic nitrogen (DON) and dissolved organic carbon (DOC) in soil. *Soil Biology & Biochemistry*, *38*, 991–999. <https://doi.org/10.1016/j.soilbio.2005.08.012>
- Kang, R. H., Mulder, J., Duan, L., & Dorsch, P. (2017). Spatial and temporal variability of soil nitric oxide emissions in N-saturated subtropical forest. *Biogeochemistry*, *134*, 337–351. <https://doi.org/10.1007/s10533-017-0368-z>
- Kaspari, M., Bujan, J., Weiser, M. D., Ning, D., Michaletz, S. T., Zhili, H., ... Wright, S. J. (2017). Biogeochemistry drives diversity in the prokaryotes, fungi, and invertebrates of a Panama forest. *Ecology*, *98*, 2019–2028. <https://doi.org/10.1002/ecy.1895>
- Kellner, H., Luis, P., Schlitt, B., & Buscot, F. (2009). Temporal changes in diversity and expression patterns of fungal laccase genes within the organic horizon of a brown forest soil. *Soil Biology & Biochemistry*, *41*, 1380–1389. <https://doi.org/10.1016/j.soilbio.2009.03.012>
- Kou, Y., Li, J., Wang, Y., Li, C., Tu, B., Yao, M., & Li, X. (2017). Scale-dependent key drivers controlling methane oxidation potential in Chinese grassland soils. *Soil Biology & Biochemistry*, *111*, 104–114. <https://doi.org/10.1016/j.soilbio.2017.04.005>
- Kuzyakov, Y. (2002). Review: Factors affecting rhizosphere priming effects. *Journal of Plant Nutrition and Soil Science*, *165*(4), 382–396. [https://doi.org/10.1002/1522-2624\(200208\)165:4<382:AID-JPLN382>3.0.CO;2-#](https://doi.org/10.1002/1522-2624(200208)165:4<382:AID-JPLN382>3.0.CO;2-#)
- Kuzyakov, Y. (2006). Sources of CO₂ efflux from soil and review of partitioning methods. *Soil Biology & Biochemistry*, *38*, 425–448. <https://doi.org/10.1016/j.soilbio.2005.08.020>
- Lammel, D. R., Feigl, B. J., Cerri, C. C., & Nusslein, K. (2015). Specific microbial gene abundances and soil parameters contribute to C, N, and greenhouse gas process rates after land use change in Southern Amazonian Soils. *Frontiers in Microbiology*, *6*, 1057. <https://doi.org/10.3389/fmicb.2015.01057>
- Levy-Booth, D. J., Prescott, C. E., & Grayston, S. J. (2014). Microbial functional genes involved in nitrogen fixation, nitrification and denitrification in forest ecosystems. *Soil Biology & Biochemistry*, *75*, 11–25. <https://doi.org/10.1016/j.soilbio.2014.03.021>
- Li, J., Li, Z. A., Wang, F. M., Zou, B., Chen, Y., Zhao, J., ... Xia, H. P. (2015). Effects of nitrogen and phosphorus addition on soil microbial community in a secondary tropical forest of China. *Biology and Fertility of Soils*, *51*, 207–215. <https://doi.org/10.1007/s00374-014-0964-1>
- Li, Y., Niu, S. L., & Yu, G. R. (2016). Aggravated phosphorus limitation on biomass production under increasing nitrogen loading: A meta-analysis. *Global Change Biology*, *22*, 934–943. <https://doi.org/10.1111/gcb.13125>
- Liu, B. B., Morkved, P. T., Frostegård, A., & Bakken, L. R. (2010). Denitrification gene pools, transcription and kinetics of NO, N₂O and N₂ production as affected by soil pH. *FEMS Microbiology Ecology*, *2*, 407–417. <https://doi.org/10.1111/j.1574-6941.2010.00856.x>
- Liu, J. X., Zhou, G. Y., Xu, Z. H., Duan, H. L., Li, Y. L., & Zhang, D. Q. (2011). Photosynthesis acclimation, leaf nitrogen concentration, and growth of four tree species over 3 years in response to elevated carbon dioxide and nitrogen treatment in subtropical China. *Journal of Soils and Sediments*, *11*, 1155–1164. <https://doi.org/10.1007/s11368-011-0398-4>
- Liu, L. L., & Greaver, T. L. (2009). A review of nitrogen enrichment effects on three biogenic GHGs: The CO₂ sink may be largely offset by stimulated N₂O and CH₄ emission. *Ecology Letters*, *12*, 1103–1117. <https://doi.org/10.1111/j.1461-0248.2009.01351.x>
- Liu, X. J., Zhang, Y., Han, W. X., Tang, A. H., Shen, J. L., Cui, Z. L., ... Zhang, F. S. (2013). Enhanced nitrogen deposition over China. *Nature*, *494*, 459–462. <https://doi.org/10.1038/nature11917>
- Lu, C. Q., & Tian, H. Q. (2013). Net greenhouse gas balance in response to nitrogen enrichment: Perspectives from a coupled biogeochemical model. *Global Change Biology*, *19*, 571–588. <https://doi.org/10.1111/gcb.12049>
- Lu, L., & Jia, Z. (2013). Urease gene-containing Archaea dominate autotrophic ammonia oxidation in two acid soils. *Environmental Microbiology*, *15*, 1795–1809. <https://doi.org/10.1111/1462-2920.12071>
- Lu, X. K., Mao, Q. G., Mo, J. M., Gilliam, F. S., Zhou, G. Y., Luo, Y. Q., ... Huang, J. (2015). Divergent responses of soil buffering capacity to long-term N deposition in three typical tropical forests with different land-use history. *Environmental Science & Technology*, *49*, 4072–4080. <https://doi.org/10.1021/es5047233>
- Lu, X. K., Vitousek, P. M., Mao, Q. G., Gilliam, F. S., Luo, Y. Q., Zhou, G. Y., ... Mo, J. M. (2018). Plant acclimation to long-term high nitrogen deposition in an N-rich tropical forest. *Proceedings of the National Academy of Sciences of the United States of America*, *115*, 5187–5192. <https://doi.org/10.1073/pnas.1720777115>
- Lu, Y. H., Lueders, T., Friedrich, M. W., & Conrad, R. (2005). Detecting active methanogenic populations on rice roots using stable isotope probing. *Environmental Microbiology*, *7*, 326–336. <https://doi.org/10.1111/j.1462-2920.2005.00697.x>
- Ma, K., Conrad, R., & Lu, Y. H. (2012). Responses of methanogen mcrA genes and their transcripts to an alternate dry/wet cycle of paddy field soil. *Applied and Environmental Microbiology*, *78*, 445–454. <https://doi.org/10.1128/aem.06934-11>
- Magill, A. H., Aber, J. D., Currie, W. S., Nadelhoffer, K. J., Martin, M. E., McDowell, W. H., ... Steudler, P. (2004). Ecosystem response to 15 years of chronic nitrogen additions at the Harvard Forest LTER, Massachusetts, USA. *Forest Ecology and Management*, *196*, 7–28. <https://doi.org/10.1016/j.foreco.2004.03.033>
- Matson, P. A., McDowell, W. H., Townsend, A. R., & Vitousek, P. M. (1999). The globalization of N deposition: Ecosystem consequences in tropical environments. *Biogeochemistry*, *46*, 67–83. <https://doi.org/10.1007/BF01007574>
- Mo, J., Zhang, W., Zhu, W., Gundersen, P., Fang, Y., Li, D., & Wang, H. (2008). Nitrogen addition reduces soil respiration in a mature tropical forest in southern China. *Global Change Biology*, *14*, 403–412. <https://doi.org/10.1111/j.1365-2486.2007.01503.x>
- Morales, S. E., Cosart, T., & Holben, W. E. (2010). Bacterial gene abundances as indicators of greenhouse gas emission in soils. *The ISME Journal*, *4*, 799–808. <https://doi.org/10.1038/ismej.2010.8>
- Pajares, S., & Bohannan, B. J. M. (2016). Ecology of nitrogen fixing, nitrifying, and denitrifying microorganisms in tropical forest soils. *Frontiers in Microbiology*, *7*, 1045. <https://doi.org/10.3389/fmicb.2016.01045>
- Prosser, J. I., & Nicol, G. W. (2012). Archaeal and bacterial ammonia-oxidisers in soil: The quest for niche specialisation and differentiation. *Trends in Microbiology*, *20*, 523–531. <https://doi.org/10.1016/j.tim.2012.08.001>
- Ramette, A., & Tiedje, J. M. (2007). Multiscale responses of microbial life to spatial distance and environmental heterogeneity in a patchy ecosystem. *Proceedings of the National Academy of Sciences of the United States of America*, *104*, 2761–2766. <https://doi.org/10.1073/pnas.0610671104>
- Rousk, J., Brookes, P. C., & Baath, E. (2009). Contrasting soil pH effects on fungal and bacterial growth suggest functional redundancy in carbon mineralization. *Applied and Environmental Microbiology*, *75*, 1589–1596. <https://doi.org/10.1128/AEM.02775-08>
- Rütting, T., Huygens, D., Boeckx, P., Staelens, J., & Klemmedtsson, L. (2013). Increased fungal dominance in N₂O emission hotspots along a natural pH gradient in organic forest soil. *Biology and Fertility of Soils*, *49*, 715–721. <https://doi.org/10.1007/s00374-012-0762-6>
- Sanchez, G. (2013). *PLS path modeling with R*. Berkeley, CA: Trowchez Editions.
- Sayer, E. J., Heard, M. S., Grant, H. K., Marthews, T. R., & Tanner, E. V. J. (2011). Soil carbon release enhanced by increased tropical forest litterfall. *Nature Climate Change*, *1*, 304–307. <https://doi.org/10.1038/nclimate1190>
- Schimel, J. (2000). Rice, microbes and methane. *Nature*, *403*, 375–377. <https://doi.org/10.1038/35000325>

- Stone, M. M., Kan, J. J., & Plante, A. F. (2015). Parent material and vegetation influence bacterial community structure and nitrogen functional genes along deep tropical soil profiles at the Luquillo Critical Zone Observatory. *Soil Biology & Biochemistry*, *80*, 273–282. <https://doi.org/10.1016/j.soilbio.2014.10.019>
- Tian, H., Chen, G., Lu, C., Xu, X., Ren, W., Zhang, B., ... Wofsy, S. (2015). Global methane and nitrous oxide emissions from terrestrial ecosystems due to multiple environmental changes. *Ecosystem Health and Sustainability*, *1*, 1–20. <https://doi.org/10.1890/EHS14-0015.1>
- Treseder, K. K. (2008). Nitrogen additions and microbial biomass: A meta-analysis of ecosystem studies. *Ecology Letters*, *11*, 1111–1120. <https://doi.org/10.1111/j.1461-0248.2008.01230.x>
- Vitousek, P. M., & Matson, P. A. (1988). Nitrogen transformations in a range of tropical forest soils. *Soil Biology & Biochemistry*, *20*, 361–367. [https://doi.org/10.1016/0038-0717\(88\)90017-X](https://doi.org/10.1016/0038-0717(88)90017-X)
- Vitousek, P. M., Porder, S., Houlton, B. Z., & Chadwick, O. A. (2010). Terrestrial phosphorus limitation: Mechanisms, implications, and nitrogen-phosphorus interactions. *Ecological Applications*, *20*(1), 5–15.
- Waldrop, M. P., Zak, D. R., Blackwood, C. B., Curtis, C. D., & Tilman, D. (2006). Resource availability controls fungal diversity across a plant diversity gradient. *Ecology Letters*, *9*, 1127–1135. <https://doi.org/10.1111/j.1461-0248.2006.00965.x>
- Wang, C., Liu, D. W., & Bai, E. (2018). Decreasing soil microbial diversity is associated with decreasing microbial biomass under nitrogen addition. *Soil Biology & Biochemistry*, *120*, 126–133. <https://doi.org/10.1016/j.soilbio.2018.02.003>
- Wang, H., Mo, J., Lu, X., Xue, J., Li, J., & Fang, Y. (2009). Effects of elevated nitrogen deposition on soil microbial biomass carbon in major subtropical forests of southern China. *Frontiers of Forestry in China*, *4*, 21–27. <https://doi.org/10.1007/s11461-009-0013-7>
- Wang, J. S., Wu, L., Zhang, C. Y., Zhao, X. H., Bu, W. S., & Gadow, K. V. (2016). Combined effects of nitrogen addition and organic matter manipulation on soil respiration in a Chinese pine forest. *Environmental Science and Pollution Research*, *23*, 22701–22710.
- Wei, Z., Jiangming, M., Yunting, F., Xiankai, L., & Hui, W. (2008). Effects of nitrogen deposition on the greenhouse gas fluxes from forest soils. *Acta Ecologica Sinica*, *28*, 2309–2319. [https://doi.org/10.1016/S1872-2032\(08\)60047-5](https://doi.org/10.1016/S1872-2032(08)60047-5)
- Yang, Y. F., Wu, L. W., Lin, Q. Y., Yuan, M. T., Xu, D. P., Yu, H., ... Zhou, J. Z. (2013). Responses of the functional structure of soil microbial community to livestock grazing in the Tibetan alpine grassland. *Global Change Biology*, *19*, 637–648. <https://doi.org/10.1111/gcb.12065>
- Yu, Y. J., Zhang, J. B., Chen, W. W., Zhong, W. H., Zhu, T. B., & Cai, Z. C. (2014). Effect of land use on the denitrification, abundance of denitrifiers, and total nitrogen gas production in the subtropical region of China. *Biology and Fertility of Soils*, *50*, 105–113. <https://doi.org/10.1007/s00374-013-0839-x>
- Zak, R., Freedma, Z. B., Upchurch, R. A., Steffens, M., & Kögel-Knabner, I. (2017). Anthropogenic N deposition increases soil organic matter accumulation without altering its biochemical composition. *Global Change Biology*, *23*, 933–944.
- Zamanian, K., Zarebanadkouki, M., & Kuzyakov, Y. (2018). Nitrogen fertilization raises CO₂ efflux from soil inorganic carbon: A global assessment. *Global Change Biology*, *24*, 2810–2817. <https://doi.org/10.1111/gcb.14148>
- Zhang, J. B., Cai, Z. C., Cheng, Y., & Zhu, T. B. (2009). Denitrification and total nitrogen gas production from forest soils of Eastern China. *Soil Biology & Biochemistry*, *41*, 2551–2557. <https://doi.org/10.1016/j.soilbio.2009.09.016>
- Zhang, J. B., Yu, Y. J., Zhu, T. B., & Cai, Z. C. (2014). The mechanisms governing low denitrification capacity and high nitrogen oxide gas emissions in subtropical forest soils in China. *Journal of Geophysical Research-Biogeosciences*, *119*, 1670–1683. <https://doi.org/10.1002/2014jg002662>
- Zhang, T., Chen, H. Y. H., & Ruan, H. (2018). Global negative effects of nitrogen deposition on soil microbes. *The ISME Journal*, *12*, 1817–1825. <https://doi.org/10.1038/s41396-018-0096-y>
- Zhang, W., Mo, J., Zhou, G., Gundersen, P., Fang, Y., Lu, X., ... Dong, S. (2008). Methane uptake responses to nitrogen deposition in three tropical forests in southern China. *Journal of Geophysical Research: Atmospheres*, *113*, D11. <https://doi.org/10.1029/2007jd009195>
- Zheng, J., Wang, Y., Hui, N., Dong, H., Chen, C. R., Han, S. J., & Xu, Z. H. (2017). Changes in CH₄ production during different stages of litter decomposition under inundation and N addition. *Journal of Soil and Sediments*, *17*(4), 949–959. <https://doi.org/10.1007/s11368-016-1628-6>
- Zheng, X. H., Fu, C. B., Xu, X. K., Yan, X. D., Huang, Y., Han, S. H., ... Chen, G. X. (2002). The Asian nitrogen cycle case study. *Ambio*, *31*, 79–87. <https://doi.org/10.1579/0044-7447-31.2.79>
- Zhou, G. Y., Liu, S. G., Li, Z., Zhang, D. Q., Tang, X. L., Zhou, C. Y., ... Mo, J. M. (2006). Old-growth forests can accumulate carbon in soils. *Science*, *314*, 1417–1417. <https://doi.org/10.1126/science.1130168>
- Zhou, J. Z., Xue, K., Xie, J. P., Deng, Y., Wu, L. Y., Cheng, X. H., ... Luo, Y. Q. (2012). Microbial mediation of carbon-cycle feedbacks to climate warming. *Nature Climate Change*, *2*, 106–110. <https://doi.org/10.1038/nclimate1331>
- Zhou, Z. H., Wang, C. K., Zheng, M. H., Jiang, L. F., & Luo, Y. Q. (2017). Patterns and mechanisms of responses by soil microbial communities to nitrogen addition. *Soil Biology & Biochemistry*, *115*, 433–441. <https://doi.org/10.1016/j.soilbio.2017.09.015>
- Zhu, B., Gutknecht, J. L. M., Herman, D. J., Keck, D. C., Firestone, M. K., & Cheng, W. X. (2014). Rhizosphere priming effects on soil carbon and nitrogen mineralization. *Soil Biology & Biochemistry*, *76*, 183–192. <https://doi.org/10.1016/j.soilbio.2014.04.033>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: Tian J, Dungait JAJ, Lu X, et al.

Long-term nitrogen addition modifies microbial composition and functions for slow carbon cycling and increased sequestration in tropical forest soil. *Glob Change Biol*. 2019;25:3267–3281. <https://doi.org/10.1111/gcb.14750>